

CRITICAL-EXPONENT SOBOLEV NORMS AND THE SLICE THEOREM FOR THE QUOTIENT SPACE OF CONNECTIONS

PAUL M. N. FEEHAN

1. INTRODUCTION

The use of certain ‘critical-exponent’ Sobolev norms is an important feature of methods employed by Taubes to solve the anti-self-dual and related non-linear elliptic partial differential equations [49, 50, 51]. Indeed, the estimates one can obtain using these critical-exponent norms appear to be the best possible when one needs to bound the norm of a Green’s operator for a Laplacian, depending on a connection varying in a non-compact family, in terms of minimal data such as the first positive eigenvalue of the Laplacian or the L^2 norm of the curvature of the connection. Despite their utility, particularly in applications where an optimal analysis is required for gluing or degeneration problems (for example, when considering degenerating families of anti-self-dual connections or stable, holomorphic vector bundles — see Section 1.4 below), these methods are not widely known. Following Taubes [48, 49, 50, 51] we describe a collection of critical-exponent Sobolev norms and general Green’s operator estimates depending only on first positive eigenvalues or the L^2 norm of the connection’s curvature. These estimates are especially useful both for the construction of gluing maps, in the case of either anti-self-dual connections [50] or, more recently, in the case of PU(2) monopoles [19, 20], and for analyzing their asymptotic behavior with respect to Uhlenbeck limits of the underlying gluing data. We apply them here to prove an optimal slice theorem for the quotient space of connections. The result is ‘optimal’ in the sense that if a point $[A]$ in the quotient space is known to be just L_1^2 -close enough to a reference point $[A_0]$ (see below for the precise statement), then A can be placed in Coulomb gauge relative to A_0 , with all constants depending at most on the first positive eigenvalue of the covariant Laplacian defined by A_0 and the L^2 norm of the curvature of A_0 . Such slice theorems are particularly advantageous when analyzing gluing maps and their differentials in situations (such as those of [21, 34] and [19]) where the underlying gluing data is allowed to ‘bubble’. In this paper we shall for simplicity only consider connections over four-dimensional manifolds, but the methods and results can be adapted to the case of manifolds of arbitrary dimension, as in [54], to prove slice theorems applicable to cases where the reference connection is allowed to degenerate.

1.1. Critical-exponent Sobolev norms and the slice theorem. Suppose that X is a closed, Riemannian four-manifold, that G is a compact Lie group, and that $\mathcal{B}_E^{k,p} = \mathcal{A}_E^{k,p} / \mathcal{G}_E^{k+1,p}$ is the quotient space of L_k^p connections on a G bundle E modulo the Banach Lie group of L_{k+1}^p gauge transformations. Here, the integer $k \geq 1$ and the Sobolev exponent $1 < p < \infty$ obey the constraint $(k+1)p > 4$, so $L_{k+1}^p(X) \subset C^0(X)$ and gauge

Date: Completed October 2, 1996. dg-ga/9711004. This version: February 1, 1999.

The author was supported in part by an NSF Mathematical Sciences Postdoctoral Fellowship under grant DMS 9306061.

transformations in $\mathcal{G}_E^{k+1,p}$ are continuous. When $(k+1)p = 4$ we have the ‘borderline’, ‘critical’, or ‘limiting case’ of the Sobolev embedding theorem: $L_{k+1}^p(X) \subset L^q(X)$ for all $q < \infty$ but not $q = \infty$.

A connection $A \in \mathcal{A}_E^{k,p}$ is in *Coulomb gauge* relative to a reference connection A_0 if $d_{A_0}^*(A - A_0) = 0$ and it is a standard result that $\mathbf{S}_{A_0} = A_0 + \text{Ker } d_{A_0}^* \subset \mathcal{A}_E^{k,p}$ provides a *slice* for the action of the gauge group $\mathcal{G}_E^{k+1,p}$ [2, 14, 22, 28, 31, 32, 39, 43]. More exactly, if $\mathbf{B}_{A_0}^{k,p}(\varepsilon)$ is the L_k^p ball in \mathbf{S}_{A_0} with center A_0 and L_{k,A_0}^p -radius ε and $\text{Stab}_{A_0} \subset \mathcal{G}_E^{k+1,p}$ is the stabilizer of A_0 , then the projection $\pi : \mathbf{B}_{A_0}^{k,p}(\varepsilon) / \text{Stab}_{A_0} \rightarrow \mathcal{B}_E^{k,p}$ is a homeomorphism onto its image and thus contains a small enough L_k^p ball

$$B_{[A_0]}^{k,p}(\eta) = \{[A] \in \mathcal{B}_E^k : \text{dist}_{L_{k,A_0}^p}([A], [A_0]) < \eta\},$$

where the gauge-invariant distance function on the quotient is defined by

$$\text{dist}_{L_{k,A_0}^p}([A], [A_0]) = \inf_{u \in \mathcal{G}_E^{k+1,p}} \|u(A) - A_0\|_{L_{k+1,A_0}^p}.$$

One unsatisfactory aspect of the standard slice theorem concerns the dependence of the constants $\varepsilon([A_0], k, p)$ and $\eta([A_0], k, p)$ above on the orbit $[A_0]$ — in particular on the curvature F_{A_0} — when k and p are large enough that gauge transformations in $\mathcal{G}_E^{k+1,p}$ are continuous. Even in the minimal cases, $k = 1$ and $p > 2$ or $k = 2$ and $p = 2$, the constants ε, η depend unfavorably on $[A_0]$ when the curvature F_{A_0} ‘bubbles’ and $[A_0]$ approaches an ideal point in the Uhlenbeck compactification \bar{M}_E of the moduli space M_E of anti-self-dual connections (that is, a point in $\bar{M}_E \setminus M_E$). This makes it difficult to analyze the asymptotic behavior of Taubes’ gluing maps [45, 47, 49, 50] and their differentials on neighborhoods of ideal points in \bar{M}_E , since the balls $\mathbf{B}_{A_0}^{k,p}(\varepsilon)$ and $B_{[A_0]}^{k,p}(\eta)$ tend to shrink as $[A_0]$ approaches a point in $\bar{M}_E \setminus M_E$. For example, if the connection A_0 is anti-self-dual, then its *energy* is bounded by a constant depending only on the topology of E via the Chern-Weil identity

$$-\frac{1}{4\pi^2} \int_X \text{tr}(F_{A_0} \wedge F_{A_0}) = p_1(\mathfrak{g}_E),$$

whereas $\|F_{A_0}\|_{L^p}$ (with $p > 2$) or $\|F_{A_0}\|_{L_{1,A_0}^2}$ tends to infinity as the curvature of A_0 becomes concentrated and $[A_0]$ approaches the Uhlenbeck boundary.

Our main purpose in this article is to prove a global analogue, Theorem 1.1, of Uhlenbeck’s local Coulomb gauge-fixing theorem [54, Theorems 1.3 & 2.1] and a corresponding slice theorem, Theorem 1.2, where the radii of the coordinate balls on the quotient $\mathcal{B}_E^{k,p}$ depend only on $\|F_{A_0}\|_{L^2}$ and the *least positive eigenvalue* $\nu_0[A_0]$ of the Laplacian $d_{A_0}^* d_{A_0}$ on $\Omega^0(\mathfrak{g}_E)$. The key difficulty in establishing Theorem 1.1 is to ensure that the constants depend at most on $\|F_{A_0}\|_{L^2}$ and $\nu_0[A_0]$: To guarantee this minimal dependence, we employ critical-exponent Sobolev norms (defined below) to circumvent the fact that when $(k+1)p = 4$ the standard Sobolev embedding and multiplication theorems fall just short of what one needs to give the quotient $\mathcal{B}_E^{k,p} = \mathcal{A}_E^{k,p} / \mathcal{G}_E^{k+1,p}$ a manifold structure (see Section 4). Such norms were introduced by Taubes for related purposes in [49].

1.2. Statement of results. For clarity, we now fix $p = 2$ and $k \geq 2$ and define the following distance functions on the quotient space $\mathcal{B}_E^k = \mathcal{A}_E^k / \mathcal{G}_E^{k+1}$ of L_k^2 connections modulo L_{k+1}^2

gauge transformations,

$$\begin{aligned} \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) &= \inf_{u \in \mathcal{G}_E^{k+1}} (\|u(A) - A_0\|_{L^{2\sharp,4}} + \|d_{A_0}^*(u(A) - A_0)\|_{L^{\sharp,2}}), \\ \text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) &= \inf_{u \in \mathcal{G}_E^{k+1}} \left(\|u(A) - A_0\|_{L_{1,A_0}^2} + \|d_{A_0}^*(u(A) - A_0)\|_{L^{\sharp,2}} \right), \end{aligned}$$

where the Sobolev norms are defined by:

$$\begin{aligned} \|a\|_{L^\sharp(X)} &= \sup_{x \in X} \|\text{dist}^{-2}(x, \cdot)|a|\|_{L^1(X)}, \\ \|a\|_{L^{\sharp,2}(X)} &= \|a\|_{L^2(X)} + \|a\|_{L^\sharp(X)}, \\ \|a\|_{L^{2\sharp}(X)} &= \sup_{x \in X} \|\text{dist}^{-1}(x, \cdot)|a|\|_{L^2(X)}, \\ \|a\|_{L^{2\sharp,4}(X)} &= \|a\|_{L^4(X)} + \|a\|_{L^{2\sharp}(X)}, \\ \|a\|_{L_{1,A_0}^2(X)} &= \left(\|a\|_{L^2(X)}^2 + \|\nabla_{A_0} a\|_{L^2(X)}^2 \right)^{1/2}, \end{aligned}$$

for any $a \in \Omega^1(\mathfrak{g}_E)$; here, $\text{dist}(x, y)$ denotes the geodesic distance between points $x, y \in X$. Like the L^4 norm, the $L^{2\sharp}$ norm on one-forms is scale-invariant. Our first result is the following global analogue of Uhlenbeck's theorem and complements results of Taubes in [49, §6]:

Theorem 1.1. *Let X be a closed, smooth four-manifold with metric g and let G be a compact Lie group. Then there are positive constants c, z with the following significance. Let E be a G bundle over X and suppose that $k \geq 2$ is an integer. Given a point $[A_0]$ in \mathcal{B}_E^k , let $\nu_0[A_0]$ be the least positive eigenvalue of the Laplacian $d_{A_0}^* d_{A_0}$ on $\Omega^0(\mathfrak{g}_E)$ and set $K_0 = (1 + \nu_0[A_0]^{-1})(1 + \|F_{A_0}\|_{L^2})$. Let ε_1 be a constant satisfying $0 < \varepsilon_1 \leq zK_0^{-2}(1 + \nu_0[A_0]^{-1/2})^{-1}$. Then the following hold:*

1. *For any $[A] \in \mathcal{B}_E^k$ with $\text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) < \varepsilon_1$, there is a gauge transformation $u \in \mathcal{G}_E^{k+1}$, unique up to an element of the stabilizer $\text{Stab}_{A_0} \subset \mathcal{G}_E^{k+1}$, such that*
 - (a) $d_{A_0}^*(u(A) - A_0) = 0$,
 - (b) $\|u(A) - A_0\|_{L^{2\sharp,4}} \leq cK_0 \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0])$.
2. *For any $[A] \in \mathcal{B}_E^k$ with $\text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) < \varepsilon_1$, there is a gauge transformation $u \in \mathcal{G}_E^{k+1}$, unique up to an element of the stabilizer $\text{Stab}_{A_0} \subset \mathcal{G}_E^{k+1}$, such that*
 - (a) $d_{A_0}^*(u(A) - A_0) = 0$,
 - (b) $\|u(A) - A_0\|_{L^{2\sharp,4}} \leq cK_0 \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0])$,
 - (c) $\|u(A) - A_0\|_{L_{1,A_0}^2} \leq cK_0 \text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0])$.

Theorem 1.1 is especially well-adapted to establishing the existence of transformations to Coulomb gauge when the point $[A] \in \mathcal{B}_E^k$ has the form $A = A_0 + d_{A_0}^{+,*}v$, with A_0 approximately anti-self-dual (so $F_{A_0}^+$ is small) and $v \in L_{k+1}^2(\Lambda^+ \otimes \mathfrak{g}_E)$, since $d_{A_0}^* d_{A_0}^{+,*}v = (F_{A_0}^+)^*v$. Points of this form in the moduli space of anti-self-dual connections, M_E , are constructed by Taubes' gluing maps [45, 47, 50].

The distance function $\text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0])$ is bounded by scale invariant norms,

$$\|a\|_{L^4(X,g)} + \|\nabla_A^g a\|_{L^2(X,g)} + \sup_{x \in X} \|\text{dist}_g^{-2}(x, \cdot)|d_{A_0}^{*g} a|\|_{L^1(X,g)}, \quad a \in \Omega^1(\mathfrak{g}_E),$$

since the $L^{4/\ell}$ norm on $\otimes^\ell(T^*X)$ is conformally invariant, while the third term is invariant under constant rescalings $g \mapsto \tilde{g} = \lambda^{-2}g$ of the metric, as $d_{A_0}^{*\tilde{g}}a = \lambda^2 d_{A_0}^{*g}a$, $\text{dist}_{\tilde{g}}^{-2}(x, y) = \lambda^2 \text{dist}_g^{-2}(x, y)$ and $dV_{\tilde{g}} = \lambda^{-4}dV_g$. Similarly for $\text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0])$.

In Theorem 2.1 of [54] the L^2 norm of the curvature F_A of a local connection matrix A over the unit ball in \mathbb{R}^4 provides a natural (gauge-invariant) measure of the distance from $[A]$ to $[\Gamma]$, where Γ is the product connection. Uhlenbeck's theorem guarantees the existence of an L_{k+1}^p gauge transformation u taking an L_k^p connection A on the product bundle over the unit four-ball, with product connection Γ , to a connection $u(A)$ satisfying $d_\Gamma^*(u(A) - \Gamma) = 0$ and $\|u(A) - \Gamma\|_{L_1^2} \leq c\|F_A\|_{L^2}$; one only requires that $\|F_A\|_{L^2}$ be smaller than a universal constant.

We next have the following refinement of the standard slice theorem for the quotient space \mathcal{B}_E^k . The observation that an L^4 -ball in $\text{Ker } d_{A_0}^*$ provides a slice for \mathcal{G}_E^{k+1} was pointed out to us Mrowka; that slightly smaller $L^{2\sharp,4}$ and L_{1,A_0}^2 balls provide slices follows from the second of our two proofs of Theorem 1.1 in Section 8. For any $\varepsilon > 0$, define

$$\begin{aligned} B_{[A_0]}^{1,\sharp,2}(\varepsilon) &= \{[A] \in \mathcal{B}_E^k : \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) < \varepsilon\} \subset \mathcal{B}_E^k, \\ B_{[A_0]}^{1,*,2}(\varepsilon) &= \{[A] \in \mathcal{B}_E^k : \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) < \varepsilon\} \subset \mathcal{B}_E^k, \\ \mathbf{B}_{A_0}^4(\varepsilon) &= \{A \in \mathcal{A}_E^k : d_{A_0}^*(A - A_0) = 0 \text{ and } \|A - A_0\|_{L^4(X)} < \varepsilon\} \subset \mathbf{S}_{A_0}, \end{aligned}$$

where $\mathbf{S}_{A_0} = \{A_0\} + \text{Ker}(d_{A_0}^*|_{L_k^2}) \subset \mathcal{A}_E^k$ is the slice through A_0 .

Theorem 1.2. *Let X be a closed, smooth four-manifold with metric g and let G be a compact Lie group. Then there are positive constants c_1, c_2, z with the following significance. Let E be a G bundle over X , let $k \geq 2$ be an integer, and suppose that $[A_0] \in \mathcal{B}_E^k$. Then the following hold:*

1. *For any constant ε_0 satisfying $0 < \varepsilon_0 < z(1 + \nu_0[A_0]^{-1/2})^{-1}$, the projection $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0)/\text{Stab}_{A_0} \rightarrow \mathcal{B}_E^k$ given by $A \mapsto [A]$ is a homeomorphism onto an open neighborhood of $[A_0] \in \mathcal{B}_E^k$ and a diffeomorphism on the open subset where $\text{Stab}_{A_0}/\text{Center}(G)$ acts freely;*
2. *For any constant ε_1 satisfying $0 < \varepsilon_1 \leq zK_0^{-2}(1 + \nu_0[A_0]^{-1/2})^{-1}$ we have the following inclusions of open neighborhoods in \mathcal{B}_E^k :*

$$B_{[A_0]}^{1,\sharp,2}(\varepsilon_1) \subset B_{[A_0]}^{1,*,2}(c_1\varepsilon_1) \subset \pi(\mathbf{B}_{A_0}^4(c_2K_0\varepsilon_1)).$$

Let $\mathcal{A}_E^{*,k} \subset \mathcal{A}_E^k$ be the subspace of connections A with minimal stabilizer $\text{Stab}_A = \text{Center}(G)$ and let $\mathcal{B}_E^{*,k} = \mathcal{A}_E^{*,k}/\mathcal{G}_E^{k+1}$. It is well-known that the quotient space \mathcal{B}_E^k is Hausdorff, that the subspace $\mathcal{B}_E^{*,k} \subset \mathcal{B}_E^k$ is an open, C^∞ Banach manifold, and that the projection $\pi : \mathcal{A}_E^{*,k} \rightarrow \mathcal{B}_E^{*,k}$ is a C^∞ principal $\mathcal{G}_E^{k+1}/\text{Center}(G)$ bundle. See Proposition 3.4 for detailed statements.

That sharper versions of the standard slice theorem (as in [14, 22, 28], for example) would hold is suggested by related results of Taubes, namely [48, Lemma A.1] and [49, Lemma 6.5]: for example, they show that if u is an L_2^2 gauge transformation intertwining L_1^2 connections A_i , $i = 1, 2$, obeying a slice condition $d_{A_0}^*(A_i - A_0) = 0$ defined by an L_1^2 connection A_0 , then u is necessarily in C^0 . Moreover, transition functions relating neighborhoods of the origin in $\text{Ker}(d_{A_0}^*|_{L_1^2})$ and $\text{Ker}(d_{A_0+a}^*|_{L_1^2})$, where a is L_{1,A_0}^2 -small, are constructed in [49,

Lemma 6.5]; the constants depend only on $\|F_{A_0}\|_{L^2}$ and $\nu_0[A_0]$. (See [49, §6] for detailed statements and related results.) The proof of Theorem 1.1 makes use of methods developed in [48, 49, 50].

1.3. Outline of the proofs. Assertion (1) of Theorem 1.2 is proved in Section 3. The proof that the projection map $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0) \rightarrow \mathcal{B}_E^k$ is a local diffeomorphism away from connections with non-minimal stabilizer essentially follows Uhlenbeck's verification of 'openness' in her proof of Theorem 2.1 in [54] via the method of continuity (see Lemma 3.6). The proof that the L^4 ball $\mathbf{B}_{A_0}^4(\varepsilon_0)$ injects into the quotient (see Lemma 3.7) was suggested to us by Mrowka. The remainder of our article is taken up with the proof of Theorem 1.1 and hence Assertion (2) of Theorem 1.2.

In Section 4 we introduce the family of critical-exponent Sobolev norms, $L_{k,A_0}^{\sharp,2}$, $k = 0, 1, 2$, used to complete the proof of Theorem 1.1 and in Section 5 we describe the crucial embedding theorems enjoyed by those Sobolev spaces, as well as estimates for the Green's operator of the Laplacian $d_{A_0}^* d_{A_0}$. In particular, $L_{k,A_0}^p \subset L_{k,A_0}^{\sharp,2}$, for every $p > 2$ while, in the other direction, $L_{2,A_0}^{\sharp,2} \subset C^0$. The latter embedding is the key motivation for the definition of these norms and it greatly facilitates the derivation of Green's operator estimates, in a wide number of applications in gauge theory [49, 50], with minimal dependence on the curvature of the connection A_0 . The main ideas and embedding results in Sections 4 and 5 are due to Taubes [48, 49, 50, 51], so these sections are essentially expository. An earlier exposition from a somewhat different perspective, due to Donaldson, of Taubes' methods and some applications appears in [12]. The estimates of Section 5 are stated only in the four-dimensional case. While we might expect all of them to hold, in some form, for higher dimensions we confine our attention to dimension four as our intended applications are primarily concerned with smooth four-manifold topology. In essence, the critical-exponent norms make a virtue out of necessity of the familiar fact that while the Green's operator of the Laplacian d^*d on $C^\infty(X)$ maps $L^p(X)$ into $L^{2p/(2-p)}(X)$ for $1 < p < 2$, it does not map $L^2(X)$ into $L^\infty(X)$ [44, Chapter V]. We recall that an Orlicz space L_φ can be used to provide the 'best target space' for an embedding of $L_2^2(X)$ [1, Chapter 8]. Here, we may instead view $L_2^{\sharp,2}(X)$ as providing the 'best domain space' for an embedding into $L^\infty(X)$, since $L_2^p(X) \subset L_2^{\sharp,2}(X) \subset L^\infty(X)$ for all $p > 2$.

We give two proofs of Theorem 1.1. For our first proof, in Section 6, we essentially follow the strategy of Uhlenbeck [54] and apply the method of continuity. The difficult step here (in establishing 'openness' — see Section 6.3) is to prove that the intrinsic, gauge-invariant $\mathcal{L}_{1,A_0}^{\sharp,2}$ and $L_{1,A_0}^{\sharp,2}$ distances in the quotient \mathcal{B}_E^k bound the $L_2^{\sharp,4}$ and L_{1,A_0}^2 norms in the slice $\mathbf{S}_{A_0} \subset \mathcal{A}_E^k$: this is the point in our first proof where we use the critical-exponent estimates derived in Section 5 to control gauge transformations. The proof of 'closedness' uses a compactness argument and is given in Section 6.2.

Our second proof of Theorem 1.1 occupies Sections 7 and 8. In Section 7 we show that the exponential map $\text{Exp} : \Omega^0(\mathfrak{g}_E) \rightarrow \mathcal{G}_E$ extends to a continuous map $\text{Exp} : L_2^{\sharp,2}(\mathfrak{g}_E) \rightarrow L_2^{\sharp,2}(\mathfrak{g}_E)$ and that the resulting space of $L_2^{\sharp,2}$ -gauge transformations $\mathcal{G}_E^{2,\sharp,2}$ is a Banach Lie group. In particular, $L_2^{\sharp,2}$ -gauge transformations are *continuous* and are contained in $\mathcal{G}_E^{2,p}$ for every $p > 2$. The Sobolev multiplication and composition results for the critical-exponent norms then allow us to apply the inverse function theorem directly in Section 8, while still ensuring that all constants depend at most on $\nu_0[A_0]$ and $\|F_{A_0}\|_{L^2}$. We first use the compactness result of Section 6.1 to establish the existence of gauge transformations w in

\mathcal{G}_E^3 which minimize the $\mathcal{L}_{1,A_0}^{\sharp,2}$ and $L_{1,A_0}^{\sharp,2}$ distances in the quotient \mathcal{B}_E^k . Then, assuming the norm $\|w(A) - A_0\|_{L^{2\sharp,4}}$ or $\|w(A) - A_0\|_{L_{1,A_0}^2}$ is sufficiently small, we use the Sobolev embedding and multiplication theorems of Sections 4, 5, and 7 and a quantitative version of the inverse function theorem to prove the existence of a gauge transformation $v \in \mathcal{G}_E^3$ such that $d_{A_0}^*(u(A) - A_0) = 0$, $u = vw \in \mathcal{G}_E^{k+1}$, and $\|u(A) - A_0\|_{L^{2\sharp,4}}$ and $\|u(A) - A_0\|_{L_{1,A_0}^2}$ are controlled by $\text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0])$ and $\text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0])$, respectively.

1.4. Applications and extensions. Applications of the methods and results of the present article occur in situations where connections, metrics, or holomorphic structures are allowed to degenerate and uniform estimates are required for elliptic operators whose coefficients depend on these degenerating geometric structures.

1.4.1. Degeneration of anti-self-dual connections. While the standard slice theorem is adequate for many applications in smooth four-manifold topology [14, 22, 23], one finds that it is rather less adequate for constructing gluing maps and analyzing their asymptotic behavior in sufficient generality to be useful in approaches to the Kotschick-Morgan conjecture [21, 34]. That conjecture asserts that the Donaldson invariants of a four-manifold with $b^+(X) = 1$, computed using metrics lying in different chambers of the positive cone of $H^2(X; \mathbb{R})/\mathbb{R}^*$, differ by terms depending only the homotopy type of X [25, 26].

Taubes' gluing maps can be used [34] to construct links of arbitrary lower-level reducibles in $\bar{M}_\kappa^w \setminus M_\kappa^w$, where E is now a $\text{U}(2)$ bundle and M_κ^w is the moduli space of anti-self-dual connections on $\mathfrak{su}(E)$, with $w_2(\mathfrak{su}(E)) = c_1(E) \pmod{2}$ and $\kappa = -\frac{1}{4}p_1(\mathfrak{su}(E))$. Any obvious approach to the conjecture places considerable demands on gluing theory since one must, at least in principle, be able to describe links of all ideal reducibles and not just isolated special cases.

For a lower-level reducible $[A_1, \mathbf{x}] \in M_{\kappa-\ell}^w \times \text{Sym}^\ell(X)$, the Laplacian $d_{A_1}^* d_{A_1}$ has a kernel of dimension equal to that of Stab_{A_1} , so the Laplacian $d_A^* d_A$ of connection A necessarily has $\dim \text{Stab}_{A_1}$ 'small eigenvalues' tending to zero as $[A]$ approaches $[A_1, \mathbf{x}]$ and the remaining eigenvalues of $d_A^* d_A$ are bounded below by one-half the least positive eigenvalue of $d_{A_1}^* d_{A_1}$. A similar phenomenon arises in [47, 50] with the Laplacian $d_A^+ d_A^{+,*}$ on $\Omega^+(\mathfrak{g}_E)$ when the 'background' connection A_1 has $\text{Ker } d_{A_1}^+ d_{A_1}^{+,*} \neq 0$: the small eigenvalues of $d_A^+ d_A^{+,*}$ represent an obstruction to perturbing approximately anti-self-dual to anti-self-dual connections near $[A_1, \mathbf{x}]$. For such neighborhoods in \bar{M}_κ^w , Theorems 1.1 and 1.2 are easily modified by replacing $\nu_0[A_0]$ with a suitable small eigenvalue cutoff, by analogy with [47, 50], and proceeding along the lines of [49, §6, Part 3]. (Without such modifications, the coordinate balls of Theorem 1.2 shrink if $[A_0] \in M_\kappa^w$ approaches a reducible point in $\bar{M}_\kappa^w \setminus M_\kappa^w$.) To illustrate applications of the methods of Sections 4 and 5 and to point to possible generalizations of the estimates in present article, we derive some elliptic estimates for $d_A^+ + d_A^*$ in Section 5.2.

1.4.2. Degeneration of $\text{PU}(2)$ monopoles. We recall that Viktor Pidstrigach and Andrei Tyurin proposed a method [41, 42] to prove Witten's conjecture concerning the relation between the Donaldson and Seiberg-Witten invariants of smooth four-manifolds [13, 55]. Their proposal uses a moduli space of solutions to the $\text{PU}(2)$ monopole equations, which are a natural generalization of the $\text{U}(1)$ monopole equations of Seiberg and Witten and the anti-self-dual equation for $\text{SO}(3)$ connections, to construct a cobordism between links of

compact moduli spaces of $U(1)$ monopoles of Seiberg-Witten type and the moduli space of anti-self-dual connections, which appear as singularities in this larger moduli space.

The problems one encounters in attempting to implement this program have much in common with those encountered in previous attempts to prove the Kotschick-Morgan conjecture [16, 17, 18]. For the purposes of proving the general gluing theorem for $PU(2)$ monopoles (which one needs to describe the above-mentioned links) we require, among other things, a useful criterion to detect when a monopole near an ideal point in the Uhlenbeck compactification of the moduli space of $PU(2)$ monopoles lies in the image of a gluing map. An L^2_{k,A_0} (with $k \geq 2$) or even an L^p_{1,A_0} (with $p > 2$) measure of closeness to the image of the approximate gluing map is not useful for this purpose as the radius of such balls tend to shrink as the curvature of the connection bubbles, as in the case of anti-self-connections, and simpler implicit-function theorem arguments used to show that a point lies in the image of a gluing map would fail in general. In [19] we use analogues of Theorems 1.1 and 1.2 and their methods of proof to facilitate arguments that the $PU(2)$ monopole gluing maps are ‘surjective’ (in the above sense) and are diffeomorphisms onto their images. The surjectivity property of Taubes’ gluing maps for anti-self-dual connections is a special case of a more general gluing result for critical points of the Yang-Mills functional [49, Proposition 8.2]. The gluing maps for anti-self-dual connections constructed by Donaldson and employed in [9, 10, 11, 14, 15] are shown to be surjective in [9, 14, 15].

1.4.3. Degeneration of stable, holomorphic vector bundles. Via work of Donaldson [7, 8] one may view the degeneration of stable holomorphic vector bundles as a special case of the degeneration of anti-self-dual connections. Comparisons of these degenerations have been given by Li [29] and Morgan [33] in their work relating the the Gieseker and Uhlenbeck compactifications of the moduli space of anti-self-dual connections over a complex, Kähler surface. The notion of stability is broadened in [4, 35, 53] by considering the vortex and nonabelian monopole equations over complex, Kähler manifolds. In particular, as explained to the author by Daskalopoulos [6], there are potential applications for the types of estimates considered here when applying methods from gauge theory to study the degeneration of holomorphic vector bundles near the boundary of Teichmüller space; see [5] for a description of the moduli space of stable, holomorphic vector bundles over Riemann surfaces. Such gauge-theoretic methods are in turn expected to have applications to three-manifold topology [6].

Acknowledgements. We warmly thank Tom Mrowka for helpful and stimulating discussions on gauge theory and his suggestion that L^4 -balls inject into the quotient, and warmly thank Cliff Taubes for very helpful discussions and especially for explaining his ideas and methods in [48, 49]. We would also like to thank the Harvard Mathematics Department and the National Science Foundation for their generous hospitality and support during the preparation of this article.

2. PRELIMINARIES

We assume throughout this article that X is a closed, connected, smooth, four-manifold with Riemannian metric g . Let G be a compact Lie group with matrix representation $\rho : G \subset SO(\mathbb{E}) = SO(r)$ where $\mathbb{E} \simeq \mathbb{R}^r$ as a real inner product space, let P be a principal G bundle, and let $E = P \times_{\rho} \mathbb{E}$ be the corresponding Riemannian vector bundle associated to P by the representation ρ . Let $\mathfrak{g}_E \subset \mathfrak{gl}(E)$ be the bundle of Lie algebras associated to P

via the adjoint representation $\text{Ad} : G \rightarrow \text{Aut}(\mathfrak{g})$ of G on its Lie algebra \mathfrak{g} and viewed as a subbundle of $\mathfrak{gl}(E)$ via the induced representation $\rho_* : \mathfrak{g} \subset \mathfrak{so}(\mathbb{R})$.

Given the covariant derivative $\nabla_A : C^\infty(E) \rightarrow C^\infty(T^*X \otimes E)$, we define the exterior covariant derivative $d_A : \Omega^i(E) \rightarrow \Omega^{i+1}(E)$ in the usual way by setting $d_A = \nabla_A$ on $\Omega^0(E) = C^\infty(E)$ and extending d_A to $\Omega^i(E) = C^\infty(\Lambda^i \otimes E)$, where $\Lambda^i := \Lambda^i(T^*X)$, according to the rule $d_A(\omega \wedge v) = d\omega \wedge v + (-1)^i \omega \wedge d_A v$ for $\omega \in \Omega^i(X)$ and $v \in \Omega^j(E)$.

For any integer $k \geq 0$, exponent $1 \leq p \leq \infty$, and L_k^p connection A_0 on E we define the L_k^p Sobolev completion, $L_k^p(\Lambda^\ell \otimes E)$, of $\Omega^\ell(E)$ with respect to the norm

$$\|s\|_{L_{k,A_0}^p(X)} := \left(\sum_{j=0}^k \|\nabla_{A_0}^j s\|_{L^p(X)}^p \right)^{1/p}.$$

We define the action of a C^∞ gauge transformation $u \in \mathcal{G}_E$ on a C^∞ connection A on the bundle E by pushforward, so $u(A) := A - (d_A u)u^{-1}$. Fix a connection $A_0 \in \mathcal{A}_E$, let $\mathcal{A}_E^k = A_0 + L_k^2(\Lambda^1 \otimes \mathfrak{g}_E)$, and define

$$\mathcal{G}_E^{k+1} := \{u \in L_{k+1}^2(\mathfrak{gl}(E)) : u \in G \text{ a.e.}\} \subset L_{k+1}^2(\mathfrak{gl}(E)).$$

The space \mathcal{G}_E^{k+1} is a Banach Lie group, with Lie algebra $T_{\text{id}_E} \mathcal{G}_E^{k+1} = L_{k+1}^2(\mathfrak{g}_E)$, and acts smoothly on \mathcal{A}_E^k with quotient $\mathcal{B}_E^k := \mathcal{A}_E^k / \mathcal{G}_E^{k+1}$ endowed with the quotient L_k^2 topology.

The stabilizer subgroup $\text{Stab}_A \subset \mathcal{G}_E^{k+1}$ for a connection A on E always contains the center $Z(G) \subset G$. We let $\mathcal{A}_E^{*,k} \subset \mathcal{A}_E^k$ denote the space of connections $A \in \mathcal{A}_E^k$ with minimal stabilizer $\text{Stab}_A = Z(G)$ and let $\mathcal{B}_E^{*,k} = \mathcal{A}_E^{*,k} / \mathcal{G}_E^{k+1}$. As usual, the stabilizer subgroup $\text{Stab}_A \subset \mathcal{G}_E$ can be identified with a closed subgroup of $G \subset \text{GL}(E|_{x_0})$ for any point $x_0 \in X$ by parallel translation with respect to the connection A . Let \mathfrak{stab}_A denote the Lie algebra of Stab_A , so $\mathfrak{stab}_A = \text{Ker}\{d_A : L_{k+1}^2(\mathfrak{g}_E) \rightarrow L_k^2(\Lambda^1 \otimes \mathfrak{g}_E)\}$.

Throughout the article, we use c or z to denote positive constants which depend at most on the Riemannian manifold (X, g) and the group G ; constants may increase from one line to the next and are not renamed unless clarity demands otherwise.

3. THE SLICE THEOREM

In this section we prove the first assertion of Theorem 1.2 — see Proposition 3.4 below — namely, that a small enough L^4 -ball $\mathbf{B}_{A_0}^4(\varepsilon_0) / \text{Stab}_{A_0}$ provides a slice for the action of \mathcal{G}_E^{k+1} . The proof that the projection $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0) / \text{Stab}_{A_0} \rightarrow \mathcal{B}_E^k$ is injective (Lemma 3.7) was suggested to us by Mrowka.

Let $k \geq 2$ be an integer. The Banach Lie group \mathcal{G}_E^{k+1} has Lie algebra $T_{\text{id}_E} \mathcal{G}_E^{k+1} = L_{k+1}^2(\mathfrak{g}_E)$ and exponential map $\text{Exp} : L_{k+1}^2(\mathfrak{g}_E) \rightarrow \mathcal{G}_E^{k+1}$ given by $\zeta \mapsto u = \text{Exp} \zeta$. Recall that $\text{Stab}_A = \{\gamma \in \mathcal{G}_E^{k+1} : \gamma(A) = A\}$ may be identified with a Lie subgroup of G and has Lie algebra $\mathfrak{stab}_A = \text{Ker}(d_A|_{L_{k+1}^2})$. The operator $d_A^* : L_{k+2}^2(\Lambda^1 \otimes \mathfrak{g}_E) \rightarrow L_{k+1}^2(\mathfrak{g}_E)$ has closed range and we have an L^2 -orthogonal decomposition

$$\begin{aligned} (3.1) \quad T_{\text{id}_E} \mathcal{G}_E^{k+1} &= L_{k+1}^2(\mathfrak{g}_E) \\ &= (\text{Ker}(d_A|_{L_{k+1}^2}))^\perp \oplus \text{Ker}(d_A|_{L_{k+1}^2}) \\ &= \text{Im}(d_A^*|_{L_{k+2}^2}) \oplus \text{Ker}(d_A|_{L_{k+1}^2}) \\ &= (\text{Ker}(d_A|_{L_{k+1}^2}))^\perp \oplus \mathfrak{stab}_A. \end{aligned}$$

Let $\text{Stab}_A^\perp = L_{k+1}^2 \cap \text{Stab}_A^\perp = \text{Exp}((\text{Ker } d_A|_{L_{k+1}^2})^\perp)$, the second equality following from the Sobolev composition lemma. The subspace $\text{Stab}_A^\perp \subset \mathcal{G}_E^{k+1}$ is closed and is a Banach submanifold of \mathcal{G}_E^{k+1} with codimension $\dim \mathfrak{stab}_A$. From Claim 3.5 below we see that Stab_A^\perp is a slice near $\text{id}_E \in \mathcal{G}_E^{k+1}$ for the right action of Stab_A on \mathcal{G}_E^{k+1} .

The map $d_A : L_{k+1}^2(\mathfrak{g}_E) \rightarrow L_k^2(\Lambda^1 \otimes \mathfrak{g}_E)$ has closed range and so we have an L^2 -orthogonal decomposition

$$\begin{aligned} (3.2) \quad T_A \mathcal{A}_E^k &= L_k^2(\Lambda^1 \otimes \mathfrak{g}_E) \\ &= \text{Im}(d_A|_{L_{k+1}^2}) \oplus \text{Ker}(d_A^*|_{L_k^2}) \\ &= \text{Im}(d_A|_{L_{k+1}^2}) \oplus \mathbf{K}_A, \end{aligned}$$

of the tangent space to the space of L_k^2 connections at the point A , where we set where $\mathbf{K}_A = \text{Ker}(d_A^*|_{L_k^2})$.

The *slice* $\mathbf{S}_A \subset \mathcal{A}_E^k$ through a connection A is given by $\mathbf{S}_A = A + \mathbf{K}_A$. If π is the projection from \mathcal{A}_E^k onto $\mathcal{B}_E^k = \mathcal{A}_E^k / \mathcal{G}_E^{k+1}$, denoted by $A \mapsto [A]$, we let

$$\begin{aligned} \mathbf{B}_A(\varepsilon) &= \{A_1 \in \mathbf{S}_A : \|A_1 - A\|_{L_{k;A}^2} < \varepsilon\} \\ &= A + \{a \in \mathbf{K}_A : \|a\|_{L_{k;A}^2} < \varepsilon\} \end{aligned}$$

be the open L_k^2 -ball in \mathbf{S}_A with center A and $L_{k,A}^2$ -radius ε . Similarly, we let

$$\begin{aligned} \mathbf{B}_A^4(\varepsilon) &= \{A_1 \in \mathbf{S}_A : \|A_1 - A\|_{L^4} < \varepsilon\} \\ &= A + \{a \in \mathbf{K}_A : \|a\|_{L^4} < \varepsilon\} \end{aligned}$$

be the open ball in \mathbf{S}_A with center A and L^4 -radius ε .

The proof that the quotient space \mathcal{A}_E^k is Hausdorff makes use of the following well-known technical result [22, Proposition A.5]. Note that the space \mathcal{G}_E^2 is neither a Banach Lie group nor does it act smoothly on \mathcal{A}_E^k for $k \geq 1$.

Lemma 3.1. *Let E be a Hermitian bundle over a Riemannian manifold X and let $k \geq 2$ be an integer. Suppose $\{A_\alpha\}$ and $\{B_\alpha\}$ are sequences of L_k^2 unitary connections on E and that $\{u_\alpha\}$ is a sequence in \mathcal{G}_E^2 such that $u_\alpha(A_\alpha) = B_\alpha$. Then the following hold.*

1. *The sequence $\{u_\alpha\}$ is in \mathcal{G}_E^{k+1} ;*
2. *If $\{A_\alpha\}$ and $\{B_\alpha\}$ converge in \mathcal{A}_E^k to limits A_∞ , B_∞ , then there is a subsequence $\{\alpha'\} \subset \{\alpha\}$ such that $\{u_{\alpha'}\}$ converges in \mathcal{G}_E^{k+1} to u_∞ and $B_\infty = u_\infty(A_\infty)$.*

We shall need the following quantitative version of the inverse function theorem here and especially in Section 8:

Theorem 3.2. *Let $\Phi : \mathbf{E} \rightarrow \mathbf{F}$ be a C^ℓ map of Banach spaces, for some $\ell \geq 1$, such that the differential $(D\Phi)_{x_0} : \mathbf{E} \rightarrow \mathbf{F}$ has a continuous inverse $(D\Phi)_{x_0}^{-1} : \mathbf{F} \rightarrow \mathbf{E}$ satisfying*

$$\|(D\Phi)_{x_0}^{-1}\| \leq K \quad \text{and} \quad \|(D\Phi)_x - (D\Phi)_{x_0}\| \leq \frac{1}{2}K^{-1}, \quad \text{if} \quad \|x - x_0\| \leq \delta,$$

for some positive constants K and δ . Then the following hold:

1. *The restriction of Φ to the ball $U = B^\mathbf{E}(x_0, \delta)$ is injective and $\Phi(U) = V$ is an open set in \mathbf{F} containing the ball $B^\mathbf{F}(\Phi(x_0), \delta/(2K))$;*
2. *The inverse map $\Phi^{-1} : V \rightarrow U$ is C^ℓ ;*
3. *If $x_1, x_2 \in B^\mathbf{E}(x_0, \delta)$, then $\|x_1 - x_2\| \leq 2K\|\Phi(x_1) - \Phi(x_2)\|$.*

For quantitative comparisons in this section, the following elementary fact will suffice:

Lemma 3.3. *Let \mathbf{E}, \mathbf{F} be Banach spaces and let $T \in \text{Hom}(\mathbf{E}, \mathbf{F})$ have a right (left) inverse S . If $\tilde{T} \in \text{Hom}(\mathbf{E}, \mathbf{F})$ satisfies $\|\tilde{T} - T\| < \|S\|^{-1}$, then \tilde{T} also has a right (left) inverse.*

Proof. If $S \in \text{Hom}(\mathbf{F}, \mathbf{E})$ is a right inverse for T , so $TS = \text{id}_{\mathbf{F}}$, then $\|(\tilde{T} - T)S\| \leq \|\tilde{T} - T\|\|S\| < 1$ and $\text{id}_{\mathbf{E}} + (\tilde{T} - T)S$ is an invertible element of the Banach algebra $\text{End}(\mathbf{E})$. Define $\tilde{S} = S(1 + (\tilde{T} - T)S)^{-1}$, so $\tilde{S}T = \text{id}_{\mathbf{E}}$ and \tilde{S} is a right inverse for \tilde{T} . Similarly for left inverses. \square

This consequence of the usual characterization of invertible elements of a Banach algebra will be invoked in the proof of Lemma 3.6.

Proposition 3.4. *Let X be a closed, Riemannian four-manifold. Then there is a positive constant z with the following significance. Let E be a G bundle over X . Suppose that $k \geq 2$ is an integer. Given $[A_0]$ in \mathcal{B}_E^k , let $\nu_0[A_0]$ be the least positive eigenvalue of the Laplacian $\Delta_{A_0}^0$ and let ε_0 be a constant satisfying $0 < \varepsilon_0 < z(1 + \nu_0[A_0]^{-1/2})^{-1}$. Then the following hold:*

1. *The space \mathcal{B}_E^k is Hausdorff;*
2. *The subspace $\mathcal{B}_E^{*,k} \subset \mathcal{B}_E^k$ is open and is a C^∞ Banach manifold with local parametrizations given by $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0) \rightarrow \mathcal{B}_E^{*,k}$;*
3. *The projection $\pi : \mathcal{A}_E^{*,k} \rightarrow \mathcal{B}_E^{*,k}$ is a C^∞ principal $\mathcal{G}_E^{k+1}/\text{Center}(G)$ bundle;*
4. *The projection $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0)/\text{Stab}_{A_0} \rightarrow \mathcal{B}_E^k$ is a homeomorphism onto an open neighborhood of $[A_0] \in \mathcal{B}_E^k$ and a diffeomorphism on the and a diffeomorphism on the open subset where $\text{Stab}_{A_0}/\text{Center}(G)$ acts freely.*

Proof. The stabilizer Stab_{A_0} acts freely on \mathcal{G}_E^{k+1} and thus on the Banach manifold $\mathcal{G}_E^{k+1} \times \mathbf{S}_{A_0}^k$ by $(u, A) \mapsto \gamma \cdot (u, A) = (u\gamma^{-1}, \gamma(A))$ and so the quotient $\mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{S}_{A_0}$ is again a Banach manifold. We define a smooth map

$$(3.3) \quad \Psi : \mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{S}_{A_0} \rightarrow \mathcal{A}_E^k, \quad [u, A] \mapsto u(A).$$

Our main task is to show that the map Ψ is (i) a local diffeomorphism onto its image and (ii) injective upon restriction to a sufficiently small neighborhood $\mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0)$. Given $\delta_0 > 0$, let $B_{\text{id}_E}(\delta_0)$ be the ball $\{u \in \mathcal{G}_E^{k+1} : \|u - \text{id}_E\|_{L_{k+1;A_0}^2} < \delta_0\}$ and let $B_{\text{id}_E}^\perp(\delta_0) = B_{\text{id}_E}(\delta_0) \cap \text{Stab}_{A_0}^\perp$.

Claim 3.5. *For small enough $\delta = \delta(A_0, k)$, the ball $B_{\text{id}_E}(\delta)$ is diffeomorphic to an open neighborhood in $B_{\text{id}_E}^\perp(\delta) \times \text{Stab}_{A_0}$, with inverse map given by $(u_0, \gamma) \mapsto u = u_0\gamma$.*

Proof. The differential of the multiplication map

$$\text{Stab}_{A_0}^\perp \times \text{Stab}_{A_0} \rightarrow \mathcal{G}_E^{k+1}, \quad (u_0, \gamma) \mapsto u_0\gamma,$$

at $(\text{id}_E, \text{id}_E)$ is given by

$$\text{Ker}(d_{A_0}|_{L_{k+1}^2})^\perp \oplus \mathfrak{stab}_{A_0} \rightarrow L_{k+1}^2(\mathfrak{g}_E) \quad (\zeta, \chi) \mapsto u_0\zeta\gamma + u_0\gamma\chi,$$

and so is just the identity map with respect to the L^2 -orthogonal decomposition (3.1) of the range. Hence, the Banach space implicit function theorem implies that there is a diffeomorphism from an open neighborhood of $(\text{id}_E, \text{id}_E)$ onto an open neighborhood of

$\text{id}_E \in \mathcal{G}_E^{k+1}$. For small enough δ , we may suppose that if $u \in B_{\text{id}_E}(\delta)$, then u can be written uniquely as $u = u_0\gamma$ with $u_0 \in B_{\text{id}_E}^\perp(\delta)$ and $\gamma \in \text{Stab}_{A_0}$. \square

Lemma 3.6. *For any $0 < \varepsilon_0 < \frac{1}{2}(1 + \nu_0[A_0]^{-1/2})^{-1}$, the map Ψ is a local diffeomorphism from $\mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0)$ onto its image in \mathcal{A}_E^k .*

Proof. We first restrict the map Ψ to a neighborhood $B_{\text{id}_E}(\delta_0) \times_{\text{Stab}_{A_0}} \mathbf{S}_{A_0}$, which is diffeomorphic to the neighborhood $B_{\text{id}_E}^\perp(\delta) \times \mathbf{S}_{A_0}$ in $\text{Stab}_{A_0}^\perp \times \mathbf{S}_{A_0}$ by Claim 3.5. The differential of the induced map

$$(3.4) \quad \Psi : \text{Stab}_{A_0}^\perp \times \mathbf{S}_{A_0} \rightarrow \mathcal{A}_E^k, \quad (u, A) \mapsto u(A),$$

at $(\text{id}_E, A) := (\text{id}_E, A_0 + a_0)$ is given by

$$(D\Psi)_{(\text{id}_E, A)} : T_{\text{id}_E} \text{Stab}_{A_0}^\perp \oplus T_A \mathbf{S}_{A_0} \rightarrow T_A \mathcal{A}_E^k, \\ (\zeta, a) \mapsto -d_A \zeta + a = -d_{A_0} \zeta - [a_0, \zeta] + a,$$

where we recall that $T_A \mathbf{S}_{A_0} = \mathbf{K}_{A_0} = \text{Ker}(d_{A_0}^*|_{L_k^2})$ and

$$T_{\text{id}_E} \text{Stab}_{A_0}^\perp = (\text{Ker}(d_{A_0}|_{L_{k+1}^2}))^\perp = \text{Im}(d_{A_0}^*|_{L_{k+2}^2}).$$

Using the L^2 -orthogonal decomposition (3.2) of the range we see that the map

$$-d_{A_0} \oplus \text{id}_E : (\text{Ker}(d_{A_0}|_{L_1^2}))^\perp \oplus \text{Ker}(d_{A_0}^*|_{L_1^2}) \rightarrow \text{Im}(d_{A_0}|_{L_1^2}) \oplus \text{Ker}(d_{A_0}^*|_{L_1^2})$$

given by $(\zeta, b) \mapsto -d_{A_0} \zeta + b$ is a Hilbert space isomorphism. More explicitly, the operator

$$d_{A_0} : (\text{Ker}(d_{A_0}|_{L_1^2}))^\perp \rightarrow \text{Im}(d_{A_0}|_{L_1^2}) = (\text{Ker}(d_{A_0}^*|_{L^2}))^\perp$$

has a two-sided inverse

$$G_{A_0}^0 d_{A_0}^* : \text{Im}(d_{A_0}|_{L_1^2}) \rightarrow (\text{Ker}(d_{A_0}|_{L_1^2}))^\perp,$$

where $G_{A_0}^0$ is the Green's operator for the Laplacian $\Delta_{A_0}^0 = d_{A_0}^* d_{A_0}$: indeed, $G_{A_0}^0 d_{A_0}^* d_{A_0} = G_{A_0}^0 \Delta_{A_0}^0$ is the L^2 -orthogonal projection $\Pi_{A_0}^0$ from $L_1^2(\Lambda^1 \otimes \mathfrak{g}_E)$ onto $(\text{Ker}(d_{A_0}|_{L_1^2}))^\perp$ and $d_{A_0} G_{A_0}^0 d_{A_0}^*$ is the L^2 -orthogonal projection $\Pi_{A_0}^{1,\perp} = \text{id} - \Pi_{A_0}^1$ from $L^2(\Lambda^1 \otimes \mathfrak{g}_E)$ onto $(\text{Ker}(d_{A_0}^*|_{L^2}))^\perp$, as

$$d_{A_0}^* (\text{id} - d_{A_0} G_{A_0}^0 d_{A_0}^*) = 0.$$

For $\zeta \in (\text{Ker}(d_{A_0}|_{L_1^2}))^\perp$ and $b = d_{A_0} \zeta \in \text{Im}(d_{A_0}|_{L_1^2})$, we have

$$\|G_{A_0}^0 d_{A_0}^* b\|_{L_{1,A_0}^2} = \|G_{A_0}^0 \Delta_{A_0}^0 \zeta\|_{L_{1,A_0}^2} = \|\Pi_{A_0}^0 \zeta\|_{L_{1,A_0}^2} = \|\zeta\|_{L_{1,A_0}^2} \\ \leq \|d_{A_0} \zeta\|_{L^2} + \|\zeta\|_{L^2} \leq (1 + \nu_0^{-1/2}) \|d_{A_0} \zeta\|_{L^2} = (1 + \nu_0^{-1/2}) \|b\|_{L^2}$$

and so $G_{A_0}^0 d_{A_0}^*$ has $\text{Hom}(L^2, L_{1,A_0}^2)$ operator norm bound

$$\|G_{A_0}^0 d_{A_0}^*\| \leq 1 + \nu_0^{-1/2}.$$

The Sobolev embedding $L_1^2 \subset L^4$ and Kato's inequality imply that

$$\|d_A \zeta - d_{A_0} \zeta\|_{L^2} \leq \|[a_0, \zeta]\|_{L^2} \leq 2\|a_0\|_{L^4} \|\zeta\|_{L^4} \leq 2\|a_0\|_{L^4} \|\zeta\|_{L_{1,A_0}^2},$$

and so $d_A - d_{A_0}$ has $\text{Hom}(L_{1,A_0}^2, L^2)$ operator norm bound

$$\|d_A - d_{A_0}\| \leq 2\|a_0\|_{L^4}.$$

In particular, we see that $(D\Psi)_{(\text{id}_E, A_0)}^{-1} = G_{A_0}^0 d_{A_0}^* \oplus \text{id} = G_{A_0}^0 d_{A_0}^* \oplus G_{A_0}^0 \Delta_{A_0}^0$ satisfies

$$\|(D\Psi)_{(\text{id}_E, A_0)}^{-1}\| \leq 1 + \nu_0^{-1/2} \quad \text{and} \quad \|(D\Psi)_{(\text{id}_E, A)} - (D\Psi)_{(\text{id}_E, A_0)}\| \leq 2\|a_0\|_{L^4}.$$

Hence, Lemma 3.3 implies that if $\|a_0\|_{L^4} < \frac{1}{2}(1 + \nu_0^{-1/2})^{-1}$, then the operator

$$(D\Psi)_{(\text{id}_E, A)} : (\text{Ker}(d_{A_0}|_{L_1^2}))^\perp \times \text{Ker}(d_{A_0}^*|_{L_1^2}) \rightarrow L^2(\Lambda^1 \otimes \mathfrak{g}_E)$$

is an isomorphism from L_1^2 to L^2 and restricts to a bounded linear map from L_{k+1}^2 to L_k^2 . Provided $(D\Psi)_{(\text{id}_E, A)} : L_{k+1}^2 \rightarrow L_k^2$ is bijective, the open mapping theorem guarantees the existence of a bounded inverse $(D\Psi)_{(\text{id}_E, A)}^{-1} : L_k^2 \rightarrow L_{k+1}^2$. If $(D\Psi)_{(\text{id}_E, A)}(\zeta, a) = 0$ for $(\zeta, a) \in L_{k+1}^2$, then (ζ, a) is zero in L_1^2 and thus zero in L_{k+1}^2 , so $(D\Psi)_{(\text{id}_E, A)}$ is injective. If $b \in L_k^2(\Lambda^1 \otimes \mathfrak{g}_E)$, then $b = (D\Psi)_{(\text{id}_E, A)}(\zeta, a) = -d_A \zeta + a$ for some $(\zeta, a) \in (\text{Ker}(d_{A_0}|_{L_1^2}))^\perp \times \text{Ker}(d_{A_0}^*|_{L_1^2})$. As $d_{A_0}^* a = 0$, we have

$$d_{A_0}^* d_A \zeta = -d_{A_0}^* b \in L_{k-1}^2$$

and $d_{A_0}^* d_A : L_{k+1}^2 \rightarrow L_{k-1}^2$ is an elliptic operator with L_{k-1}^2 coefficients. Thus, $\zeta \in L_{k+1}^2$, so $a = b + d_A \zeta \in L_k^2$, and $(D\Psi)_{(\text{id}_E, A)}$ is surjective.

Combining the above observations, we see that the operator

$$(D\Psi)_{(\text{id}_E, A)} : (\text{Ker}(d_{A_0}|_{L_{k+1}^2}))^\perp \oplus \text{Ker}(d_{A_0}^*|_{L_k^2}) \rightarrow L_k^2(\Lambda^1 \otimes \mathfrak{g}_E),$$

is an isomorphism for all $A = A_0 + a_0$ with $\|a_0\|_{L^4} < \varepsilon_0 = \frac{1}{2}(1 + \nu_0^{-1/2})^{-1}$. So, by the Banach space implicit function theorem, there are positive constants $\varepsilon = \varepsilon(A, k)$ and $\delta = \delta(A, k)$ and an open neighborhood $U_A \subset \mathcal{A}_E^k$ such that the map

$$\Psi : B_{\text{id}_E}^\perp(\delta) \times \mathbf{B}_A(\varepsilon) \rightarrow U_A, \quad (u, A_1) \mapsto u(A_1),$$

with $\mathbf{B}_A(\varepsilon) \subset \mathbf{B}_{A_0}^4(\varepsilon_0)$, gives a diffeomorphism from an open neighborhood of (id_E, A) onto an open neighborhood of A . In particular, we obtain a map $U_A \rightarrow \text{Stab}_{A_0}^\perp$, given by $A_1 \mapsto u = u_{A_1}$, such that

$$\Psi^{-1}(A_1) = (u, u^{-1}(A_1)) \in B_{\text{id}_E}^\perp(\delta) \times \mathbf{B}_A(\varepsilon) \subset \text{Stab}_{A_0}^\perp \times \mathbf{B}_{A_0}^4(\varepsilon_0).$$

Hence, for any $A_1 \in U_A$ there is a unique $u \in B_{\text{id}_E}^\perp(\delta)$ such that $u^{-1}(A_1) - A_0 \in \mathbf{K}_{A_0}$:

$$(3.5) \quad d_{A_0}^*(u^{-1}(A_1) - A_0) = 0.$$

The neighborhood $\mathbf{B}_{A_0}^4(\varepsilon_0)$ is Stab_{A_0} -invariant: if $A \in \mathbf{B}_{A_0}(\varepsilon)$ and $\gamma \in \text{Stab}_{A_0}$, then

$$\|\gamma(A) - A_0\|_{L^4} = \|A - \gamma^{-1}(A_0)\|_{L^4} = \|A - A_0\|_{L^4} < \varepsilon,$$

and

$$\begin{aligned} d_{A_0}^*(\gamma(A) - A_0) &= \gamma \left(d_{\gamma^{-1}(A_0)}^*(A - \gamma^{-1}(A_0)) \right) \\ &= \gamma(d_{A_0}^*(A - A_0)) = 0, \end{aligned}$$

so $\gamma(A) \in \mathbf{B}_{A_0}(\varepsilon)$.

The group \mathcal{G}_E^{k+1} acts on $\mathcal{G}_E^{k+1} \times \mathbf{S}_{A_0}$ by $(u, A) \mapsto (vu, A)$, and so gives a diffeomorphism

$$B_{\text{id}_E}(\delta) \times \mathbf{B}_{A_0}^4(\varepsilon_0) \rightarrow B_v(\delta) \times \mathbf{B}_{A_0}^4(\varepsilon_0), \quad (u, A) \mapsto (vu, A),$$

and as this action commutes with the given action of Stab_{A_0} , it descends to a diffeomorphism

$$B_{\text{id}_E}(\delta) \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0) \rightarrow B_v(\delta) \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0), \quad [u, A] \mapsto [vu, A],$$

for each $v \in \mathcal{G}_E^{k+1}$. Consequently, the \mathcal{G}_E^{k+1} -equivariant map

$$\mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0) \rightarrow \mathcal{A}_E^k$$

is a local diffeomorphism onto its image, as desired. \square

Plainly, $[\gamma(A)] = [A]$ for each $\gamma \in \text{Stab}_{A_0}$ and $A \in \mathbf{B}_{A_0}^4(\varepsilon_0)$ and hence, the projection $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0) \rightarrow \mathcal{A}_E^k$ factors through $\mathbf{B}_{A_0}^4(\varepsilon)/\text{Stab}_{A_0}$.

Lemma 3.7. *There is a positive constant z with the following significance. Let $\nu_0[A_0]$ be the least positive eigenvalue of the Laplacian $\Delta_{A_0}^0$. Then for any constant ε_0 satisfying $0 < \varepsilon_0 < z(1 + \nu_0[A_0]^{-1/2})^{-1}$, the projection map $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0)/\text{Stab}_{A_0} \rightarrow \mathcal{B}_E^k$ is injective.*

Proof. Suppose $A_i \in \mathbf{B}_{A_0}^4(\varepsilon_0)$ for $i = 1, 2$ and that $[A_1] = [A_2] \in \mathcal{B}_E^k$, so $u(A_1) = A_2$ for some $u \in \mathcal{G}_E^{k+1}$. Since $u(A_0) = A_0 - (d_{A_0}u)u^{-1}$, we see that $u \in \text{Stab}_{A_0}$ if and only if $d_{A_0}u = 0$. Here, we view $u \in L_{k+1}^2(\mathfrak{gl}(E))$ via the isometric embedding $\mathcal{G}_E^{k+1} \subset L_{k+1}^2(\mathfrak{gl}(E))$ and write

$$u = u_0 - \gamma,$$

where $u_0 \in (\text{Ker } d_{A_0})^\perp$ and $\gamma \in \text{Ker } d_{A_0}$. We claim that $u_0 = 0$, so $u = \gamma \in \text{Stab}_{A_0}$.

Since $u(A_1) := A_1 - (d_{A_1}u)u^{-1} = A_2$, we have $A_2u = A_1u - d_{A_1}u = A_1u - d_{A_0}u - [A_1 - A_0, u]$, and therefore

$$d_{A_0}u_0 = d_{A_0}u = u(A_1 - A_0) - (A_2 - A_0)u.$$

Since $d_{A_0}^*(A_i - A_0) = 0$ for $i = 1, 2$, we obtain

$$\begin{aligned} d_{A_0}^*d_{A_0}u_0 &= - * (d_{A_0}u \wedge *(A_1 - A_0)) + ud_{A_0}^*(A_1 - A_0) \\ &\quad - (d_{A_0}^*(A_2 - A_0))u + *(*(A_2 - A_0) \wedge d_{A_0}u) \\ &= - * (d_{A_0}u_0 \wedge *(A_1 - A_0)) + *(*(A_2 - A_0) \wedge d_{A_0}u_0). \end{aligned}$$

Integrating by parts gives

$$\|d_{A_0}u_0\|_{L^2}^2 = (d_{A_0}^*d_{A_0}u_0, u_0)_2 \leq \|d_{A_0}^*d_{A_0}u_0\|_{L^{4/3}}\|u_0\|_{L^4}.$$

Kato's inequality and the embedding $L_1^2 \subset L^4$ gives $\|u_0\|_{L^4} \leq c(\|d_{A_0}u_0\|_{L^2} + \|u_0\|_{L^2})$, so the eigenvalue estimate $\|u_0\|_{L^2} \leq \nu_0^{-1/2}\|d_{A_0}u_0\|_{L^2}$ gives $\|u_0\|_{L^4} \leq c(1 + \nu_0^{-1/2})\|d_{A_0}u_0\|_{L^2}$ and thus

$$\|d_{A_0}u_0\|_{L^2}^2 \leq (1 + \nu_0^{-1/2})\|d_{A_0}^*d_{A_0}u_0\|_{L^{4/3}}\|d_{A_0}u_0\|_{L^2}.$$

Therefore, if $d_{A_0}u_0 \neq 0$, the preceding expression for $d_{A_0}^*d_{A_0}u_0$ yields

$$\begin{aligned} \|d_{A_0}u_0\|_{L^2} &\leq c(1 + \nu_0^{-1/2})\|d_{A_0}^*d_{A_0}u_0\|_{L^{4/3}} \\ &\leq c(1 + \nu_0^{-1/2})\|d_{A_0}u_0\|_{L^2}(\|A_1 - A_0\|_{L^4} + \|A_2 - A_0\|_{L^4}), \end{aligned}$$

and so we have

$$1 \leq c(1 + \nu_0^{-1/2})(\|A_1 - A_0\|_{L^4} + \|A_2 - A_0\|_{L^4}) \leq c(1 + \nu_0^{-1/2})\varepsilon_0.$$

which gives a contradiction for $\varepsilon_0 < c^{-1}(1 + \nu_0^{-1/2})^{-1}$ \square

We now return to consider the local diffeomorphism Ψ of Claim 3.6. Suppose $\Psi[u_1, A_1] = \Psi[u_2, A_2] \in \mathcal{A}_E^k$, where $[u_1, A_1], [u_2, A_2] \in \mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0)$, and so $u_1(A_1) = u_2(A_2) \in \mathcal{A}_E^k$ and hence $[A_1] = [A_2] \in \mathcal{B}_E^k$. Provided ε_0 also satisfies the constraints of Claim 3.7, we have $u_2^{-1}u_1 = \gamma \in \text{Stab}_{A_0}$ and $\gamma(A_1) = A_2$. Hence $[u_2, A_2] = [u_1\gamma^{-1}, \gamma(A_1)] = [u_1, A_1]$, so Ψ is injective and therefore a diffeomorphism onto \mathcal{A}_E^k .

The map $\pi : \mathbf{B}_{A_0}^4(\varepsilon_0)/\text{Stab}_{A_0} \rightarrow \mathcal{B}_E^k$ can be factored as the composition of the inclusion $A \mapsto (\text{id}_E, A)$ of $\mathbf{B}_{A_0}^4(\varepsilon_0)$ into $\mathcal{G}_E^{k+1} \times \mathbf{B}_{A_0}^4(\varepsilon_0)$, the projection onto the Stab_{A_0} -quotient $\mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0)$, the diffeomorphism Ψ of $\mathcal{G}_E^{k+1} \times_{\text{Stab}_{A_0}} \mathbf{B}_{A_0}^4(\varepsilon_0)$ with \mathcal{A}_E^k and the projection from \mathcal{A}_E^k onto the \mathcal{G}_E^{k+1} -quotient $\mathcal{B}_E^k = \mathcal{A}_E^k/\mathcal{G}_E^{k+1}$. Hence, π is a homeomorphism onto an open neighborhood of $[A_0]$ in \mathcal{B}_E^k and a diffeomorphism on the open subset where $\text{Stab}_{A_0}/\text{Center}(G)$ acts freely.

Claim 3.8. *The quotient space \mathcal{B}_E^k is Hausdorff.*

Proof. Let Γ be the subspace $\{\{A, u(A)\} : A \in \mathcal{A}_E^k \text{ and } u \in \mathcal{G}_E^{k+1}\}$ of $\mathcal{A}_E^k \times \mathcal{A}_E^k$. If $\{(A_\alpha, u_\alpha(A_\alpha))\}$ is a sequence in Γ which converges in L_k^2 to a point $\{A_\infty, B_\infty\}$, then Lemma 3.1 implies that there is a subsequence $\{\alpha'\} \subset \{\alpha\}$ such that $\{u_{\alpha'}\}$ converges in L_{k+1}^2 to $u_\infty \in \mathcal{G}_E^{k+1}$ and $u_\infty(A_\infty) = B_\infty$. Thus, Γ is closed and the quotient $\mathcal{A}_E^k/\mathcal{G}_E^{k+1}$ is Hausdorff. \square

Claim 3.8 gives Assertion (1) of the proposition and Assertions (2), (3), and (4) now follow from the preceding arguments and Lemma 3.7. This completes the proof of the proposition. \square

4. CRITICAL-EXPONENT SOBOLEV NORMS

We now describe the basic properties of the critical-exponent norms and corresponding Banach spaces introduced by Taubes in [48, 49, 50, 51]. In particular, we give the basic embedding, multiplication, and composition lemmas we need to complete the proof of our slice theorem. We shall make frequent use of the pointwise Kato inequality, $|d|v|| \leq |\nabla_A v|$ for $v \in \Omega^0(E)$, so that the norms of the embedding and multiplication maps depend at most on the Riemannian manifold (X, g) . Moreover, for simplicity, we confine our attention to the case of closed four-manifolds: there are obvious analogues of the Sobolev lemmas described here for any n -manifold, with $n > 2$. Similarly, extensions are possible to the case of complete manifolds bounded geometry (bounded curvature and injectivity radius uniformly bounded from below) — see [1, 3] for further details for Sobolev embedding results in those situations and for the construction of Green kernels. We refer the reader to the monograph of R. Adams [1] for a comprehensive treatment of Sobolev spaces and to that of E. Stein [44] for a treatment based on potential functions.

Throughout this section, A, B denote C^∞ orthogonal connections on Riemannian vector bundles E, F over X with C^∞ sections u, v , respectively. We first have the following analogues of the L^2 and L^4 norms,

$$(4.1) \quad \begin{aligned} \|u\|_{L^\sharp(X)} &= \sup_{x \in X} \|\text{dist}^{-2}(x, \cdot)|u|\|_{L^1(X)}, \\ \|u\|_{L^{2\sharp}(X)} &= \sup_{x \in X} \|\text{dist}^{-1}(x, \cdot)|u|\|_{L^2(X)}, \end{aligned}$$

where $\text{dist}(x, y)$ denotes the geodesic distance between points x and y in X defined by the metric g ; these norms have the same behavior as the L^2 and L^4 norms with respect to constant rescalings of the metric g — the L^\sharp norm on two-forms and the $L^{2\sharp}$ norm on one-forms are *scale invariant*. Indeed, one sees this by noting that if $g \mapsto \tilde{g} = \lambda^{-2}g$, then $\text{dist}_{\tilde{g}}(x, y) = \lambda^{-1} \text{dist}_g(x, y)$ and $dV_{\tilde{g}} = \lambda^{-4}dV_g$, while for any $a \in \Omega^1(E)$ and $v \in \Omega^2(E)$, we have $|a|_{\tilde{g}} = \lambda|a|_g$, and $|v|_{\tilde{g}} = \lambda^2|v|_g$.

Next, we define analogues of the L_1^2 and L_2^2 norms

$$\begin{aligned}\|u\|_{L_{1,A}^2(X)} &= \|\nabla_A u\|_{L^2(X)} + \|u\|_{L^2(X)}, \\ \|u\|_{L_{2,A}^2(X)} &= \|\nabla_A^2 u\|_{L^2(X)} + \|\nabla_A u\|_{L^2(X)} + \|u\|_{L^2(X)},\end{aligned}$$

and set

$$\begin{aligned}(4.2) \quad \|u\|_{L_{1,A}^\sharp(X)} &= \|\nabla_A u\|_{L^\sharp(X)} + \|u\|_{L^{2^\sharp}(X)} + \|u\|_{L^\sharp(X)}, \\ \|u\|_{L_{2,A}^\sharp(X)} &= \|\nabla_A^* \nabla_A u\|_{L^\sharp(X)} + \|u\|_{L^\sharp(X)},\end{aligned}$$

where $\nabla_A^* = - * \nabla_A * : \Omega^1(E) \rightarrow \Omega^0(E)$ is the L^2 -adjoint of the map $\nabla_A : \Omega^0(E) \rightarrow \Omega^1(E)$.

Finally, we define analogues of the $C^0 \cap L_2^2$ norm

$$\|u\|_{C^0 \cap L_{2,A}^2(X)} = \|u\|_{C^0(X)} + \|u\|_{L_{2,A}^2(X)},$$

and set

$$\begin{aligned}(4.3) \quad \|u\|_{L^{\sharp,2}(X)} &= \|u\|_{L^\sharp \cap L^2(X)} = \|u\|_{L^\sharp(X)} + \|u\|_{L^2(X)}, \\ \|u\|_{L^{2^\sharp,4}(X)} &= \|u\|_{L^{2^\sharp} \cap L^4(X)} = \|u\|_{L^{2^\sharp}(X)} + \|u\|_{L^4(X)}, \\ \|u\|_{L_{1,A}^{\sharp,2}(X)} &= \|u\|_{L_{1,A}^\sharp \cap L_{1,A}^2(X)} = \|u\|_{L_{1,A}^\sharp(X)} + \|u\|_{L_{1,A}^2(X)}, \\ \|u\|_{L_{2,A}^{\sharp,2}(X)} &= \|u\|_{L_{2,A}^\sharp \cap L_{2,A}^2(X)} = \|u\|_{L_{2,A}^\sharp(X)} + \|u\|_{L_{2,A}^2(X)}.\end{aligned}$$

It might have appeared, at first glance, a little more natural to continue the obvious pattern and instead define $\|u\|_{L_{2,A}^\sharp(X)}$ using $\|\nabla_A^2 u\|_{L^\sharp(X)}$: as we shall below, though, the given definition is most useful in practice. For related reasons, if $u \in \Omega^1(E) = \Omega^0(\Lambda^1 \otimes E)$, it is convenient to define the norm $\|u\|_{L_{1,A}^\sharp(X)}$ by

$$(4.4) \quad \|u\|_{L_{1,A}^\sharp(X)} = \|\nabla_A^* u\|_{L^\sharp(X)} + \|u\|_{L^{2^\sharp}(X)} + \|u\|_{L^\sharp(X)}.$$

Let $L^\sharp(X)$ be the Banach space completion of $C^\infty(X)$ with respect to the norm $\|\cdot\|_{L^\sharp}$ and similarly define the remaining Banach spaces above.

We have the following extensions of the standard Sobolev embedding theorem [22, 37]: their proofs are given in the next section. See also [12], [40], [48], [49, §6], [50, Eq. (3.4) & §5], and [51, Lemma 4.7].

Lemma 4.1. *The following are continuous embeddings:*

1. $L_k^p(E) \subset L_k^\sharp(E)$, for $k = 0, 1, 2$ and all $p > 2$;
2. $L^q(E) \subset L^{2^\sharp}(E)$, for all $q > 4$;
3. $L_1^2(E) \subset L^{2^\sharp}(E)$.

In the reverse direction we have:

Lemma 4.2. *The following are continuous embeddings:*

1. $L^\sharp(E) \subset L^1(E)$ and $L^{2^\sharp}(E) \subset L^2(E)$;
2. $L_2^\sharp(E) \subset C^0 \cap L_1^2(E)$;

We next consider the extension of the standard Sobolev multiplication lemma [22, 37]. While there is no continuous multiplication map $L_2^2 \times L_2^2 \rightarrow L_2^2$, it is worth observing that

there is a continuous bilinear map $C^0 \cap L_2^2(E) \times C^0 \cap L_2^2(F) \rightarrow C^0 \cap L_2^2(E \otimes F)$ given by $(u, v) \mapsto u \otimes v$. Note that for $u \in \Omega^0(E)$ and $v \in \Omega^0(F)$ we have

$$(4.5) \quad \begin{aligned} \nabla_{A \otimes B}^2(u \otimes v) &= (\nabla_A^2 u) \otimes v + 2\nabla_A u \otimes \nabla_B v + u \otimes \nabla_B^2 v, \\ \nabla_{A \otimes B}^* \nabla_{A \otimes B}(u \otimes v) &= (\nabla_A^* \nabla_A u) \otimes v + *((\nabla_A u) \wedge \nabla_B v) \\ &\quad - *(\nabla_A u \wedge * \nabla_B v) + u \otimes \nabla_B^* \nabla_B v. \end{aligned}$$

Similarly, for $u \in \Omega^0(\Lambda^1 \otimes E)$ and $v \in \Omega^0(F)$, we have

$$(4.6) \quad \nabla_{A \otimes B}^*(u \otimes v) = (\nabla_A^* u) \otimes v + *(*u \wedge \nabla_B v)$$

In particular, we see that if $u, v \in \Omega^0(\mathfrak{gl}(E))$, then

$$(4.7) \quad \begin{aligned} \nabla_A^* \nabla_A(uv) &= (\nabla_A^* \nabla_A u)v + *((\nabla_A u) \wedge \nabla_A v) \\ &\quad - *(\nabla_A u \wedge (* \nabla_A v)) + u(\nabla_A^* \nabla_A v), \end{aligned}$$

an identity we will need in the next section.

Lemma 4.3. *Let $\Omega^0(E) \times \Omega^0(F) \rightarrow \Omega^0(E \otimes F)$ be given by $(u, v) \mapsto u \otimes v$. Then the following hold.*

1. *The map $C^0(E) \otimes L^\sharp(F) \rightarrow L^\sharp(E \otimes F)$ is continuous;*
2. *The map $L^{2\sharp}(E) \otimes L^{2\sharp}(F) \rightarrow L^\sharp(E \otimes F)$ is continuous;*
3. *The spaces $L_1^\sharp(F)$, $L_1^2(F)$, and $L_2^\sharp(F)$ are $L_2^\sharp(E)$ -modules;*
4. *The spaces $L_1^2(F)$, $L_1^{\sharp,2}(F)$, and $L_2^{\sharp,2}(F)$ are $L_2^{\sharp,2}(E)$ -modules;*

The conclusions continue to hold for $\Omega^1(E)$ in place of $\Omega^0(E)$ and the norms on $L_1^\sharp(\Lambda^1 \otimes E)$ and $L_1^{\sharp,2}(\Lambda^1 \otimes E)$ defined via (4.4).

Proof. Let $u \in C^\infty(E)$ and $v \in C^\infty(F)$ and denote the covariant derivatives on E , F , and $E \otimes F$ by ∇ . Using $\nabla(u \otimes v) = (\nabla u)v + u \otimes \nabla v$ and the embedding $L_2^\sharp(E) \subset C^0(E)$, we see that

$$\begin{aligned} \|u \otimes v\|_{L^\sharp} &\leq \|u\|_{C^0} \|v\|_{L^\sharp} \quad \text{and} \quad \|u \otimes v\|_{L^\sharp} \leq \|u\|_{L^{2\sharp}} \|v\|_{L^{2\sharp}}, \\ \|\nabla(u \otimes v)\|_{L^\sharp} &\leq \|\nabla u\|_{L^{2\sharp}} \|v\|_{L^{2\sharp}} + \|u\|_{C^0} \|\nabla v\|_{L^\sharp} \leq c\|u\|_{L_2^\sharp} \|v\|_{L_1^\sharp}, \\ \|\nabla(u \otimes v)\|_{L^2} &\leq \|\nabla u\|_{L^4} \|v\|_{L^4} + \|u\|_{C^0} \|\nabla v\|_{L^2} \leq \|u\|_{C^0 \cap L_2^2} \|v\|_{L_1^2}, \end{aligned}$$

and hence the multiplication maps $C^0 \times L^\sharp \rightarrow L^\sharp$, $L^{2\sharp} \times L^{2\sharp} \rightarrow L^\sharp$, and $L_2^\sharp \times L_1^\sharp \rightarrow L_1^\sharp$ are continuous. Moreover,

$$\|\nabla(u \otimes v)\|_{L^2} \leq \|\nabla u\|_{L^4} \|v\|_{L^4} + \|u\|_{C^0} \|\nabla v\|_{L^2} \leq c\|u\|_{C^0 \cap L_2^2} \|v\|_{L_1^2},$$

and so, using the embedding $L_2^\sharp \subset C^0$, the multiplication $L_2^{\sharp,2} \times L_1^{\sharp,2} \rightarrow L_1^{\sharp,2}$ is continuous. Thus, L_2^\sharp is an L_1^\sharp -module and $L_2^{\sharp,2}$ is an $L_1^{\sharp,2}$ -module.

Finally, to see that L_2^\sharp and $L_2^{\sharp,2}$ are algebras, we use the identities (4.5), noting that

$$\begin{aligned} \|\nabla^* \nabla(u \otimes v)\|_{L^\sharp} &\leq \|\nabla^* \nabla u\|_{L^\sharp} \|v\|_{C^0} + 2\|\nabla u\|_{L^{2\sharp}} \|\nabla v\|_{L^{2\sharp}} + \|u\|_{C^0} \|\nabla^* \nabla v\|_{L^\sharp} \\ &\leq c\|u\|_{L_2^\sharp} \|v\|_{L_2^\sharp}, \end{aligned}$$

so the multiplication $L_2^\sharp \times L_2^\sharp \rightarrow L_2^\sharp$ is continuous, while

$$\begin{aligned} \|\nabla^2(u \otimes v)\|_{L^2} &\leq \|\nabla^2 u\|_{L^2} \|v\|_{C^0} + 2\|\nabla u\|_{L^4} \|\nabla v\|_{L^4} + \|u\|_{C^0} \|\nabla^2 v\|_{L^2} \\ &\leq c\|u\|_{C^0 \cap L_2^2} \|v\|_{C^0 \cap L_2^2}. \end{aligned}$$

The embedding $L_2^\sharp \subset C^0$ now implies that the multiplication $L_2^{\sharp,2} \times L_2^{\sharp,2} \rightarrow L_2^{\sharp,2}$ is continuous. \square

5. CRITICAL-EXPONENT SOBOLEV EMBEDDINGS AND ESTIMATES FOR GREEN'S OPERATORS

We continue the notation and assumptions of Section 4. Our goal in this section is to prove the Sobolev embedding Lemmas 4.1 and 4.2, and to derive estimates for the Green's operator G_A of the Laplacian $\nabla_A^* \nabla_A$ on $\Omega^0(E)$. The key estimates described in this section are due to Taubes and they arise, in a variety of contexts, in [40, 48, 49, 50, 51]. However, we find it convenient to collect them here — together with some useful extensions and generalizations — both for the purposes of the present article and applications in [6, 21, 34].

5.1. Estimates for the covariant Laplacian $\nabla_A^* \nabla_A$. Let $G(x, y)$ be the kernel function for the Green's operator $(d^*d + 1)^{-1}$ of the Laplacian $d^*d + 1$ on $C^\infty(X)$. The kernel $G(x, y)$ of $(d^*d + 1)^{-1}$ behaves like $\text{dist}^{-2}(x, y)$ as $\text{dist}(x, y) \rightarrow 0$ (see [51, Lemma 4.7] and [50, §5]):

Lemma 5.1. *The kernel $G(x, y)$ is a positive C^∞ function away from the diagonal in $X \times X$ and as $\text{dist}(x, y) \rightarrow 0$,*

$$G(x, y) = \frac{1}{4\pi^2 \text{dist}^2(x, y)} + o(\text{dist}^{-2}(x, y)).$$

Proof. These and other properties of G are obtained by explicitly constructing G from an initial choice of parametrix H for $d^*d + 1$ using the method of [3, §4.2.2–3], where the kernel for the Green's operator for d^*d is constructed. Recall from [44, p. 132] that the kernel $G_0(x, y)$ for $(d^*d + 1)^{-1}$ on \mathbb{R}^4 with its standard metric satisfies

$$G_0(x, y) = \frac{1}{4\pi^2 |x - y|^2} + o(|x - y|^{-2}), \quad |x - y| \rightarrow 0.$$

The kernel G is now constructed using G_0 by following the method of [3, §4.2.2–3]. \square

Lemma 5.1 implies that there is a constant c depending at most on g such that for all $x \neq y$ in X ,

$$(5.1) \quad c^{-1} \text{dist}^{-2}(x, y) \leq G(x, y) \leq c \text{dist}^{-2}(x, y).$$

Consequently, for all $u \in \Omega^0(E)$, we have

$$(5.2) \quad c^{-1} \|u\|_{L^\sharp(X)} \leq \|G|u|\|_{C^0(X)} \leq c \|u\|_{L^\sharp(X)}.$$

Lemma 4.2 will follow from the next estimate; a similar inequality is stated as Equation (3.4) in [50] for the case $p = 2$; see [40, Lemma 5.4] for the proof when $p = 2$ and $X = \mathbb{R}^4$.

Lemma 5.2. *Let X be a closed, oriented four-manifold with metric g and let $1 \leq p < 4$. Then there is a positive constant C such that for all $f \in L_1^p(X)$,*

$$\sup_{x \in X} \|\text{dist}^{-1}(x, \cdot) f\|_{L^p} \leq C \|f\|_{L_1^p}.$$

Proof. We may assume that $f \in C^\infty(X)$. Let ϱ denote the injectivity radius of the Riemannian manifold (X, g) and let $\{B(x_i, \varrho)\}_{i=1}^N$ be a covering of X by N geodesic balls with

centers at points x_i and radius $\varrho/2$. Let $\{\chi_i\}_{i=1}^N$ be a partition of unity subordinate to this cover, so that $\text{supp } \chi_i \subset B(x_i, \varrho)$ and $\sum_{i=1}^N \chi_i = 1$. Using $f = \sum_{i=1}^N \chi_i f$, we have

$$\sup_{x \in X} \|\text{dist}^{-1}(x, \cdot) f\|_{L^p}^p \leq \sum_{i=1}^N \sup_{x \in B(x_i, \varrho)} \int_{B(x_i, \varrho)} \text{dist}^{-p}(x, \cdot) |\chi_i f|^p dV.$$

For any fixed $x \in X$, let $\exp_x^{-1} : B(x, \varrho) \rightarrow \mathbb{R}^4$ be a geodesic normal coordinate chart centered at the point $x \in X$, let $\{y^\mu\}$ be the induced local coordinates centered at $x \in X$, and let $r = \text{dist}(x, y)$. We apply the divergence theorem and integration by parts to bound the integrals

$$\int_{B(x_i, \varrho)} \text{dist}^{-p}(x, \cdot) |\chi_i f|^p dV.$$

For this purpose define vector fields $\eta_i = \sum_\mu \eta_i^\mu \partial/\partial y^\mu$ supported in $B(x_i, \varrho)$ by setting

$$\eta_i^\mu = y^\mu r^{-p} |\chi_i f|^p, \quad i = 1, \dots, N.$$

If $g = \det(g_{\mu\nu})$ then, since $r^2 = \sum_\mu (y^\mu)^2$, we have $\sum_\mu y^\mu \partial/\partial y^\mu = r \partial/\partial r$ and so

$$\begin{aligned} \text{div}_g \eta_i &= \frac{1}{\sqrt{g}} \sum_\mu \frac{\partial}{\partial y^\mu} (\eta_i^\mu \sqrt{g}) \\ &= \text{div}_{g_e} \eta_i + \sum_\mu \eta_i^\mu \frac{\partial}{\partial y^\mu} \log \sqrt{g} \\ &= \text{div}_{g_e} \eta_i + r^{1-p} |\chi_i f|^p \frac{\partial}{\partial r} \log \sqrt{g}, \end{aligned}$$

where $\text{div}_{g_e} \eta_i = \sum_\mu \partial \eta_i^\mu / \partial y^\mu$ is the divergence of η_i with respect to the pullback of the Euclidean metric g_e on $B(x_i, \varrho)$. From [3, Theorem 1.53] we recall that $\partial(\log \sqrt{g})/\partial r = O(r)$ and thus

$$\text{div}_g \eta_i = \text{div}_{g_e} \eta_i + O(r^{2-p} |\chi_i f|^p).$$

Using $\sum_\mu y^\mu \partial/\partial y^\mu = r \partial/\partial r$, we have

$$\begin{aligned} \text{div}_{g_e} \eta_i &= \sum_\mu \frac{\partial}{\partial y^\mu} (y^\mu r^{-p} |\chi_i f|^p) \\ &= (4-p) r^{-p} |\chi_i f|^p + p r^{1-p} |\chi_i f|^{p-1} \frac{\partial |\chi_i f|}{\partial r}, \end{aligned}$$

and so as $p < 4$,

$$\begin{aligned} r^{-p} |\chi_i f|^p &= \frac{1}{4-p} \text{div}_{g_e} \eta_i - \frac{p}{4-p} r^{1-p} |\chi_i f|^{p-1} \frac{\partial |\chi_i f|}{\partial r} \\ &\leq \frac{1}{4-p} \text{div}_g \eta_i + \frac{c}{4-p} r^{2-p} |\chi_i f|^p + \frac{p}{4-p} r^{1-p} |\chi_i f|^{p-1} \frac{\partial |\chi_i f|}{\partial r}. \end{aligned}$$

Note that $\partial/\partial r$ is a unit vector with respect to the metric g . The divergence theorem, Hölder's inequality, and the fact that η_i has compact support in $B(x_i, \varrho)$ imply that

$$\begin{aligned} & \int_{B(x_i, \varrho)} r^{-p} |\chi_i f|^p dV \\ & \leq \frac{p}{4-p} \int_{B(x_i, \varrho)} r^{1-p} |\chi_i f|^{p-1} |d(\chi_i f)| dV + \frac{c}{4-p} \int_{B(x_i, \varrho)} r^{2-p} |\chi_i f|^p dV \\ & \leq \frac{p}{4-p} \left(\int_{B(x_i, \varrho)} r^{-p} |\chi_i f|^p dV \right)^{(p-1)/p} \left(\int_{B(x_i, \varrho)} |d(\chi_i f)|^p dV \right)^{1/p} \\ & \quad + \frac{c}{4-p} \left(\int_{B(x_i, \varrho)} r^{-p} |\chi_i f|^p dV \right)^{(p-1)/p} \left(\int_{B(x_i, \varrho)} r^p |\chi_i f|^p dV \right)^{1/p}, \end{aligned}$$

Consequently,

$$\|\text{dist}^{-1}(x, \cdot) \chi_i f\|_{L^p} \leq C (\|d(\chi_i f)\|_{L^p} + \|\chi_i f\|_{L^p}) \leq C (\|df\|_{L^p} + \|f\|_{L^p}),$$

for each i and so the desired result follows. \square

Proof of Lemma 4.1. Define $1 \leq p' < 2$ by setting $1 = 1/p + 1/p'$. Then Hölder's inequality implies that

$$\begin{aligned} \|\text{dist}^{-2}(x, \cdot) |u|\|_{L^1} & \leq \|\text{dist}^{-2}(x, \cdot)\|_{L^{p'}} \|u\|_{L^p} \leq C \|u\|_{L^p}, \\ \|\text{dist}^{-1}(x, \cdot) |u|\|_{L^2} & \leq \|\text{dist}^{-1}(x, \cdot)\|_{L^{2p'}} \|u\|_{L^{2p}} \leq C \|u\|_{L^{2p}}, \end{aligned}$$

which gives Assertions (1) and (2). By Lemma 5.2 and Kato's inequality, $|d|u|| \leq |\nabla_A u|$, we see that

$$\begin{aligned} \sup_{x \in X} \|\text{dist}^{-1}(x, \cdot) u\|_{L^p} & = \sup_{x \in X} \|\text{dist}^{-1}(x, \cdot) |u|\|_{L^p} \\ & \leq C (\|d|u|\|_{L^p} + \|u\|_{L^p}) \\ & \leq C (\|\nabla_A u\|_{L^p} + \|u\|_{L^p}). \end{aligned}$$

Taking $p = 2$ gives Assertion (3). \square

Lemma 4.2 will follow from the estimates below; the key estimates (1) and (2) in Lemma 5.3 below and the estimates (1), (2), and (3) in Lemma 5.4 are essentially those of Lemma 6.2 in [49], except that the dependence of the constant on $\|F_A\|_{L^2}$ is made explicit, but the argument is the same as that of [49].

Lemma 5.3. *Let X be a closed, oriented four-manifold with metric g . Then there is a constant c with the following significance. Let E be a Riemannian vector bundle over X and let A be an orthogonal L_2^2 connection on E with curvature F_A . Then $L_2^\sharp(E) \subset C^0 \cap L_1^2(E)$ and the following estimates hold:*

- (1) $\|\nabla_A u\|_{L^{2^\sharp}(X)} + \|u\|_{C^0(X)} \leq c \|\nabla_A^* \nabla_A u\|_{L^\sharp(X)} + \|u\|_{L^\sharp(X)},$
- (2) $\|\nabla_A u\|_{L^{2^\sharp}(X)} + \|u\|_{C^0(X)} \leq c \|\nabla_A^* \nabla_A u\|_{L^\sharp(X)} + \|u\|_{L^2(X)},$
- (3) $\|u\|_{L^1(X)} \leq c \|u\|_{L^\sharp(X)},$
- (4) $\|u\|_{L^2(X)} \leq c \|u\|_{L^{2^\sharp}(X)},$
- (5) $\|\nabla_A u\|_{L^2(X)} \leq c \|\nabla_A u\|_{L^{2^\sharp}(X)}.$

Proof. For any $u \in C^\infty(E)$ there is the following pointwise identity [22, p. 93],

$$|\nabla_A u|^2 + \frac{1}{2} d^* d |u|^2 = \langle \nabla_A^* \nabla_A u, u \rangle,$$

and thus:

$$|\nabla_A u|^2 + \frac{1}{2} (1 + d^* d) |u|^2 = \langle \nabla_A^* \nabla_A u, u \rangle + \frac{1}{2} |u|^2,$$

Using this identity and the fact that $\int_X G(x, \cdot) (d^* d + 1) |u|^2 dV = |u|^2(x)$, we obtain

$$\int_X G(x, \cdot) |\nabla_A u|^2 dV + \frac{1}{2} |u|^2(x) \leq \int_X G(x, \cdot) |\langle \nabla_A^* \nabla_A u, u \rangle| dV + \frac{1}{2} \int_X G(x, \cdot) |u|^2 dV.$$

Therefore, from (5.1), we have

$$\begin{aligned} \|\nabla_A u\|_{L^\sharp}^2 + \| |u|^2 \|_{C^0} &\leq c \|\langle \nabla_A^* \nabla_A u, u \rangle\|_{L^\sharp} + c \| |u|^2 \|_{L^\sharp} \\ &\leq c \|\nabla_A^* \nabla_A u\|_{L^\sharp} \|u\|_{C^0} + c \|u\|_{L^\sharp} \|u\|_{C^0} \end{aligned}$$

Consequently, using rearrangement with the last term, we see that

$$\|\nabla_A u\|_{L^{2^\sharp}} + \|u\|_{C^0} \leq c \|\nabla_A^* \nabla_A u\|_{L^\sharp} + c \|u\|_{L^\sharp},$$

giving (1). Combining this estimate with the embedding and interpolation inequalities, $\|u\|_{L^\sharp} \leq c \|u\|_{L^4} \leq c \|u\|_{L^2}^{1/2} \|u\|_{C^0}^{1/2}$, and again using rearrangement with the last term yields the bound in (2). Since X is closed, for all $x \neq y$ we have $\text{dist}(x, y) \leq M < \infty$, so

$$\int_X \text{dist}^{-2}(x, \cdot) |u| dV \geq M^{-2} \int_X |u| dV,$$

and this gives the estimates in (3), (4), and (5). \square

Proof of Lemma 4.2. From Lemma 5.3 we have the estimate

$$\|u\|_{C^0} \leq c \|u\|_{L_{2, A_0}^\sharp},$$

for any $u \in C^\infty(E)$. Let $\{u_m\}$ be a sequence in $C^\infty(E)$ converging to $u \in L_2^\sharp(E)$. The sequence $\{u_m\}$ is Cauchy in $L_2^\sharp(E)$ and applying the preceding estimate to the differences $u_{m_2} - u_{m_1}$, we see that it is Cauchy in the Banach space $C^0(E)$ and so the limit u lies in $C^0(E)$. The same argument, with estimates (1) and (5) of Lemma 5.3, shows that $u \in L_1^2(E)$ and this yields Assertion (2) of the lemma. Assertion (1) follows in the same manner. \square

Lemma 5.4. *Continue the hypotheses of Lemma 5.3. Then for any $u \in (C^0 \cap L_2^2)(E)$, we have*

- (1)
$$\begin{aligned} \|\nabla_A^2 u\|_{L^2(X)} &\leq \|\nabla_A^* \nabla_A u\|_{L^2(X)} + c \|F_A\|_{L^2(X)}^{1/2} \|\nabla_A u\|_{L^4(X)} \\ &\quad + \|F_A\|_{L^2(X)} \|u\|_{C^0(X)}, \end{aligned}$$
- (2)
$$\|\nabla_A u\|_{L^4(X)} \leq \|u\|_{C^0(X)}^{1/2} (\|\nabla_A^* \nabla_A u\|_{L^2(X)} + 2 \|\nabla_A^2 u\|_{L^2(X)})^{1/2},$$
- (3)
$$\|\nabla_A^2 u\|_{L^2(X)} \leq 2 \|\nabla_A^* \nabla_A u\|_{L^2(X)} + c \|F_A\|_{L^2(X)} \|u\|_{C^0(X)}.$$

Proof. The Bochner-Weitzenböck formula for the covariant Laplacian [28, Appendix, Theorem II.1] asserts that

$$(5.3) \quad d_A^* d_A + d_A d_A^* = \nabla_A^* \nabla_A + \{F_A, \cdot\},$$

where we use $\{\cdot, \cdot\}$ to denote a certain bilinear map whose precise form is unimportant here. Integrating by parts and noting that $d_A = \nabla_A$ and $d_A^* d_A = \nabla_A^* \nabla_A$ on $\Omega^0(X, V)$ and $F_A = d_A \circ d_A$ gives

$$\begin{aligned} \|\nabla_A^2 u\|_{L^2}^2 &= (\nabla_A^* \nabla_A \nabla_A u, \nabla_A u)_{L^2} \\ &= ((d_A^* d_A + d_A d_A^*) d_A u, d_A u)_{L^2} - (\{F_A, d_A u\}, \nabla_A u)_{L^2} \\ &= (d_A^* F_A u, d_A u)_{L^2} + (d_A (d_A^* d_A) u, d_A u)_{L^2} - (\{F_A, \nabla_A u\}, \nabla_A u)_{L^2} \\ &= (F_A u, F_A u)_{L^2} + (\nabla_A^* \nabla_A u, \nabla_A^* \nabla_A u)_{L^2} - (\{F_A, \nabla_A u\}, \nabla_A u)_{L^2}. \end{aligned}$$

Therefore, applying Hölder's inequality, we find that

$$\|\nabla_A^2 u\|_{L^2}^2 \leq \|\nabla_A^* \nabla_A u\|_{L^2}^2 + c \|F_A\|_{L^2} \|\nabla_A u\|_{L^4}^2 + \|F_A\|_{L^2}^2 \|u\|_{C^0}^2,$$

and taking square roots gives the desired bound in (1).

We now use integration by parts and Kato's inequality $|d|u| \leq |\nabla_A u|$ to obtain an L^4 bound on $d_A u$:

$$\begin{aligned} \|d_A u\|_{L^4}^4 &= (d_A u, |d_A u|^2 d_A u)_{L^2} \\ &= (u, |d_A u|^2 d_A^* d_A u)_{L^2} + 2(u, |d_A u| d_A u \wedge d|d_A u|)_{L^2}, \\ &= \|u\|_{C^0} \|d_A u\|_{L^4}^2 \|d_A^* d_A u\|_{L^2} + 2\|u\|_{C^0} \|d_A u\|_{L^4}^2 \|\nabla_A d_A u\|_{L^2}, \end{aligned}$$

and so, if $d_A u \neq 0$,

$$\|d_A u\|_{L^4} \leq \|u\|_{C^0}^{1/2} (\|d_A^* d_A u\|_{L^2} + 2\|\nabla_A^2 u\|_{L^2})^{1/2},$$

which gives the desired estimate in (2).

By combining the L^4 estimate for $\nabla_A u$ with the L^2 estimate for $\nabla_A^2 u$, we obtain

$$\begin{aligned} \|\nabla_A^2 u\|_{L^2} &\leq \|\nabla_A^* \nabla_A u\|_{L^2} + \|F_A\|_{L^2} \|u\|_{C^0} \\ &\quad + c \|F_A\|_{L^2}^{1/2} \|u\|_{C^0}^{1/2} (\|\nabla_A^* \nabla_A u\|_{L^2} + \|\nabla_A^2 u\|_{L^2})^{1/2}. \end{aligned}$$

We now use rearrangement with the last term above to give

$$\|\nabla_A^2 u\|_{L^2} \leq 2\|\nabla_A^* \nabla_A u\|_{L^2} + c \|F_A\|_{L^2} \|u\|_{C^0},$$

and this establishes the desired bound in (3). \square

Lemma 5.5. *Continue the hypotheses of Lemma 5.3. Then for any $u \in L_2^{\sharp, 2}(E)$, we have:*

$$\|u\|_{L_{2,A}^2(X)} + \|u\|_{C^0(X)} \leq c(1 + \|F_A\|_{L^2(X)}) \left(\|\nabla_A^* \nabla_A u\|_{L^{\sharp, 2}(X)} + \|u\|_{L^2(X)} \right).$$

Proof. From Assertion (3) of Lemm 5.4 we have the estimate

$$\|\nabla_A^2 u\|_{L^2} \leq 2\|\nabla_A^* \nabla_A u\|_{L^2} + c \|F_A\|_{L^2} \|u\|_{C^0},$$

while integration by parts gives

$$\|\nabla_A u\|_{L^2} = (\nabla_A^* \nabla_A u, u)_{L^2}^{1/2} \leq \frac{1}{\sqrt{2}} (\|\nabla_A^* \nabla_A u\|_{L^2} + \|u\|_{L^2}).$$

According to Lemma 5.3 we have

$$\|u\|_{C^0} \leq c \|\nabla_A^* \nabla_A u\|_{L^{\sharp}} + c \|u\|_{L^2},$$

and therefore the desired bound follows by combining these estimates. \square

The above lemmas lead to the following estimates for the Green's operator $G_A : L^{\sharp,2}(E) \rightarrow L^{\sharp,2}(E)$ of the Laplacian $\nabla_A^* \nabla_A : L^{\sharp,2}(E) \rightarrow L^{\sharp,2}(E)$. For $u \in \Omega^0(E)$ define

$$(5.4) \quad \|u\|_{L^{\sharp,2}_{2,A}(X)} = \|\nabla_A^* \nabla_A u\|_{L^{\sharp,2}(X)} + \|u\|_{L^{\sharp,2}(X)},$$

and observe that this is equivalent to the $L^{\sharp,2}_{2,A}$ norm defined in Section 4, although the comparison depends on the L^2 norm of the curvature F_A .

Lemma 5.6. *Continue the hypotheses of Lemma 5.3. Let $\nu_0[A]$ be the least positive eigenvalue of the Laplacian $\nabla_A^* \nabla_A$. Then for any $u \in L^{\sharp,2} \cap (\text{Ker } \nabla_A^* \nabla_A)^\perp$, we have:*

$$\begin{aligned} (1) \quad & \|G_A u\|_{L^{\sharp,2}_{2,A}(X)} \leq c(1 + \nu_0[A]^{-1}) \|u\|_{L^{\sharp,2}(X)}, \\ (2) \quad & \|G_A u\|_{L^{\sharp,2}_{2,A}(X)} \leq c(1 + \nu_0[A]^{-1}) \|u\|_{L^{\sharp,2}(X)}, \\ (3) \quad & \|G_A u\|_{L^{\sharp,2}_{2,A}(X)} \leq c(1 + \nu_0[A]^{-1})(1 + \|F_A\|_{L^2(X)}) \|u\|_{L^{\sharp,2}(X)}. \end{aligned}$$

Proof. The first and second assertions follow from Lemma 5.3, the fact that $\nabla_A^* \nabla_A G_A u = u$ for $u \in (\text{Ker } \nabla_A^* \nabla_A)^\perp$, and the eigenvalue estimate $\|u\|_{L^2} \leq \nu_0[A]^{-1} \|\nabla_A^* \nabla_A u\|_{L^2}$, while the third assertion follows from the first and Lemma 5.5. \square

5.2. Elliptic estimates for $d_A^+ + d_A^*$. To illustrate their application and to point to possible extensions, we note that the estimates of Section 5.1 for the covariant Laplacian $\nabla_A^* \nabla_A = d_A^* d_A$ on $\Omega^0(E)$ naturally extend to give estimates for the covariant Laplacians $\nabla_A^* \nabla_A$ on $\Omega^\ell(E) = \Omega^0(\Lambda^\ell \otimes E)$. Estimates for $\nabla_A^* \nabla_A$ on $\Omega^0(\Lambda^1 \otimes E)$ and $\Omega^0(\Lambda^+ \otimes E)$ are of particular interest since these can in turn be profitably compared (via the Bochner-Weitzenböck formulas [22, Equations (6.25) & (6.26)], as in [50]) with the remaining Laplacians defined by the elliptic deformation complex for the anti-self-dual equation [14, 22], namely $d_A d_A^* + d_A^{+,*} d_A^+$ on $\Omega^0(\Lambda^1 \otimes \mathfrak{g}_E)$ and $d_A^+ d_A^{+,*}$ on $\Omega^0(\Lambda^+ \otimes \mathfrak{g}_E)$. Indeed, if B_1 and B_+ are the Levi-Civita connections on Λ^1 and Λ^+ induced by the Levi-Civita connection on TX for the metric g , then the curvature ' F_A ' in the estimates of the preceding subsection is simply replaced by [30, p. 165]

$$(5.5) \quad \begin{aligned} F_{B_1 \otimes A} &= F_{B_1} \otimes \text{id}_{\mathfrak{g}_E} + \text{id}_{\Lambda^1} \otimes F_A, \\ F_{B_+ \otimes A} &= F_{B_+} \otimes \text{id}_{\mathfrak{g}_E} + \text{id}_{\Lambda^+} \otimes F_A, \end{aligned}$$

where F_{B_1} and F_{B_+} are expressed in terms of the Riemann curvature tensor Rm and where we abuse notation slightly and denote the connections on E and \mathfrak{g}_E both by A . (See [22, Appendix C] and [28, Appendix II].) In the interests of brevity we shall confine our attention to the case of L_ℓ^p estimates with $p = 2$, though the methods can be modified to obtain estimates for $p \neq 2$ (some work is required — see [14, p. 426] for hints).

In order to compute the required elliptic estimates for d_A^+ we will need the Bochner-Weitzenböck formulas,

$$(5.6) \quad d_A d_A^* + 2d_A^* d_A^+ = \nabla_A^* \nabla_A + \{\text{Ric}, \cdot\} - 2\{F_A^-, \cdot\},$$

$$(5.7) \quad 2d_A^+ d_A^* = \nabla_A^* \nabla_A - 2\{W^+, \cdot\} + \frac{R}{3} + \{F_A^+, \cdot\},$$

for the Laplacians on $\Omega^1(\mathfrak{g}_E)$ and $\Omega^+(\mathfrak{g}_E)$ [22, Equations (6.25) & (6.26)]; here, Ric , W^+ , and R are the Ricci, self-dual Weyl, and scalar curvatures of the Riemannian metric g on X . In applications to the degeneration of anti-self-dual or 'almost anti-self-dual' connections A as in [45, 47, 49, 50], we can usually arrange to have a uniform L^∞ bound on F_A^+ , but

not a uniform L^p bound on F_A^- when $p > 2$. We derive estimates in the remainder of this subsection with such applications and assumptions in view. To illustrate the nature of the difficulty we first derive a naive $L_{1,A}^2$ estimate for $a \in L_1^2(\mathfrak{g}_E)$ in terms of the L^2 norm of $(d_A^* + d_A^+)a$:

Lemma 5.7. *Let X be a closed, oriented four-manifold with metric g . Then there is a constant c with the following significance. Let E be a Riemannian vector bundle over X and let A be an orthogonal L_A^2 connection on E with curvature F_A . Then for any $a \in L_1^2(\Lambda^1 \otimes \mathfrak{g}_E)$,*

$$(5.8) \quad \|a\|_{L_{1,A}^2(X)} \leq \sqrt{2} \|(d_A^* + d_A^+)a\|_{L^2(X)} + c(1 + \|F_A^-\|_{C^0(X)})^{1/2} \|a\|_{L^2(X)}.$$

*If a is L^2 -orthogonal to $\text{Ker } d_A^+$, so that $a = d_A^*v$ for some $v \in L_2^2(\Lambda^+ \otimes \mathfrak{g}_E)$, then*

$$(5.9) \quad \begin{aligned} \|d_A^*v\|_{L_{1,A}^2(X)} &\leq \sqrt{2} \|d_A^+ d_A^*v\|_{L^2(X)} + c(1 + \|F_A^-\|_{C^0(X)})^{1/2} \|v\|_{L^2(X)} \\ &\quad + \|F_A^+\|_{C^0(X)} \|v\|_{L^2(X)}. \end{aligned}$$

Proof. From the Bochner-Weitzenböck formula for $d_A d_A^* + 2d_A^* d_A^+$ in (5.6) and integration by parts, we have:

$$\begin{aligned} \|\nabla_A a\|_{L^2}^2 &= (\nabla_A^* \nabla_A a, a) \\ &= (d_A d_A^* a, a) + 2(d_A^* d_A^+ a, a) - (\{\text{Ric}, a\}, a) + 2(\{F_A^-, a\}, a) \\ &\leq \|d_A^* a\|_{L^2}^2 + 2\|d_A^+ a\|_{L^2}^2 + c(1 + \|F_A^-\|_{C^0}) \|a\|_{L^2}^2 \end{aligned}$$

which gives (5.8). If $a = d_A^*v$, then $d_A^* d_A^* v = (d_A^+ d_A)^* v = (F_A^+)^* v$, so that

$$\|d_A^* d_A^* v\|_{L^2} \leq \|F_A^+\|_{C^0} \|v\|_{L^2}.$$

Thus, (5.9) follows from (5.8) and the above inequality. \square

Since $d_A^* + d_A^+$ is an elliptic operator, estimates of the above form follow from the general theory of linear elliptic operators. However, the preceding elementary derivation using the Bochner-Weitzenböck formula gives us a constant whose dependence on the curvature terms F_A^- and F_A^+ is made explicit. In particular, we see that the estimate is only useful when we have a uniform C^0 bound on F_A^- independent of A , which is not possible when A bubbles. At the cost of introducing a slightly stronger norm than the L^2 norm on the right hand side of the estimate above, we can derive an $L_{1,A}^2$ bound for $a = d_A^*v$ with an estimate constant depending on $\|F_A^-\|_{L^2(X)}$ rather than $\|F_A^-\|_{C^0(X)}$. Specifically, Equation (5.5) and Lemma 5.5 give the following $L_{2,A}^2$ estimates for sections of $\Lambda^+ \otimes \mathfrak{g}_E$:

Lemma 5.8. *Continue the hypotheses of Lemma 5.7. Then the following estimate holds for any $v \in L_2^{\sharp,2}(\Lambda^+ \otimes \mathfrak{g}_E)$:*

$$\|v\|_{L_{2,A}^2(X)} + \|v\|_{C^0(X)} \leq c(1 + \|F_A\|_{L^2(X)}) (\|\nabla_A^* \nabla_A v\|_{L^{\sharp,2}(X)} + \|v\|_{L^2(X)}).$$

We now replace the covariant Laplacian $\nabla_A^* \nabla_A$ in the estimates of Lemma 5.8 by the Laplacian $d_A^+ d_A^*$ via the Bochner formula (5.7) to give:

Lemma 5.9. *Continue the hypotheses of Lemma 5.8. Then there is a positive constant $\varepsilon = \varepsilon(c)$ such that the following holds. If $\|F_A\|_{L^{\sharp,2}(X)} < \varepsilon$, then*

$$\|v\|_{L_{2,A}^2(X)} + \|v\|_{C^0(X)} \leq c(1 + \|F_A\|_{L^2(X)}) (\|d_A^+ d_A^* v\|_{L^{\sharp,2}(X)} + \|v\|_{L^2(X)}).$$

Proof. From (5.5) and Lemma 4.3 we have

$$\|\nabla_A^* \nabla_A v\|_{L^{\sharp,2}} \leq 2\|d_A^+ d_A^* v\|_{L^{\sharp,2}} + c\|v\|_{L^{\sharp,2}} + c\|F_A^+\|_{L^{\sharp,2}}\|v\|_{C^0}.$$

Combining the preceding estimate with that of Lemma 5.8, together with the embedding and interpolation inequalities $\|v\|_{L^{\sharp}} \leq c\|v\|_{L^4} \leq c\|v\|_{L^2}^{1/2}\|v\|_{C^0}^{1/2}$, and using rearrangement with the last term yields the desired bound. In particular, by choosing $\varepsilon(c)$ small enough that $c\|F_A^+\|_{L^{\sharp,2}}\|v\|_{C^0} \leq 1/2$, we may use rearrangement to bring the right-hand term $\|v\|_{C^0}$ to the left-hand side. \square

Since $\|d_A^* v\|_{L_{1,A}^2} \leq \|v\|_{L_{2,A}^2}$, Lemma 5.9 yields an $L_{1,A}^2$ estimate for $d_A^* v$:

Corollary 5.10. *Continue the hypotheses of Lemma 5.9. Then:*

$$\|d_A^* v\|_{L_{1,A}^2(X)} \leq c(1 + \|F_A\|_{L^2(X)})(\|d_A^+ d_A^* v\|_{L^{\sharp,2}(X)} + \|v\|_{L^2(X)}).$$

Note that if $a \in \Omega^1(\mathfrak{g}_E)$ is L^2 -orthogonal to $\text{Ker } d_A^+$, so that $a = d_A^* v$ for some $v \in \Omega^+(\mathfrak{g}_E)$, and $\text{Ker } d_A^+ d_A^{+,*} = 0$, then the estimate of Corollary 5.10 can be written in the more familiar form

$$(5.10) \quad \|a\|_{L_{1,A}^2(X)} \leq c(1 + \|F_A\|_{L^2(X)})(\|d_A^+ a\|_{L^{\sharp,2}(X)} + \nu_2[A]^{-1/2}\|a\|_{L^2(X)}),$$

where we make use of the eigenvalue estimate $\|v\|_{L^2} \leq \nu_2[A]^{-1/2}\|d_A^* v\|_{L^2}$; the term $d_A^+ a$ above can be replaced by $(d_A^+ + d_A^*)a$ without changing the estimate constants. Here, $\nu_2[A]$ is the least positive eigenvalue of the Laplacian $d_A^+ d_A^{+,*}$.

6. EXISTENCE OF GAUGE TRANSFORMATIONS VIA THE METHOD OF CONTINUITY

In this section we complete the proof of Theorem 1.1 using the method of continuity. The strategy broadly follows that of Uhlenbeck's proof of Theorem 2.1 in [54]. The main new technical difficulty, not present in [54], is the need to compare distances in the Coulomb-gauge slice $\mathbf{S}_{A_0} \subset \mathcal{A}_E^k$ through the connection A_0 and gauge-invariant distances in \mathcal{B}_E^k from the point $[A_0]$. It is at this stage of the method of continuity (in proving openness — see Lemma 6.6) — that we need to employ the special norms and Green's operator estimates described in Sections 4 and 5 in order to achieve the requisite C^0 control of gauge transformations; the proof of closedness works, as one would expect, with standard Sobolev L^4 and L_1^2 norms. In [54], the L^2 norm of the curvature F_A essentially serves as a gauge-invariant L_1^2 measure of distance from $[A]$ to $[\Gamma]$, where Γ is the product connection on the product G bundle over the unit ball.

For $k \geq 2$, we define the following distance functions on \mathcal{B}_E^k ,

$$\begin{aligned} \text{dist}_{L^4}([A], [A_0]) &= \inf_{u \in \mathcal{G}_E^{k+1}} \|u(A) - A_0\|_4, \\ \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) &= \inf_{u \in \mathcal{G}_E^{k+1}} (\|u(A) - A_0\|_{L^{\sharp,4}} + \|d_{A_0}^*(u(A) - A_0)\|_{L^{\sharp,2}}), \\ \text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) &= \inf_{u \in \mathcal{G}_E^{k+1}} \left(\|u(A) - A_0\|_{L_{1,A_0}^2} + \|d_{A_0}^*(u(A) - A_0)\|_{L^{\sharp,2}} \right), \\ \text{dist}_{L_{\ell,A_0}^2}([A], [A_0]) &= \inf_{u \in \mathcal{G}_E^{k+1}} \|u(A) - A_0\|_{L_{\ell,A_0}^2}, \quad 1 \leq \ell \leq k, \end{aligned}$$

and the following balls with center at $[A_0]$ and radius ε :

$$\begin{aligned}\bar{B}_{[A_0]}^{1,*,2}(\varepsilon) &= \{[A] \in \mathcal{B}_E^k : \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) \leq \varepsilon\}, \\ \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon) &= \{[A] \in \mathcal{B}_E^k : \text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) \leq \varepsilon\}, \\ \bar{B}_{[A_0]}^{\ell,2}(\varepsilon) &= \{[A] \in \mathcal{B}_E^k : \text{dist}_{L_{\ell,A_0}^2}([A], [A_0]) \leq \varepsilon\}.\end{aligned}$$

Their open counterparts are denoted $B_{[A_0]}^{1,*,2}(\varepsilon)$, $B_{[A_0]}^{1,\sharp,2}(\varepsilon)$, and $B_{[A_0]}^{\ell,2}(\varepsilon)$, respectively. Our goal in this section is to establish the following:

Theorem 6.1. *Let X be a closed, smooth four-manifold with metric g and let G be a compact Lie group. Then there are positive constants c, z with the following significance. Let E be a G bundle over X and suppose that $k \geq 2$ is an integer. Given a point $[A_0]$ in \mathcal{B}_E^k , let $\nu_0[A_0]$ be the least positive eigenvalue of the Laplacian $\nabla_{A_0}^* \nabla_{A_0}$ on $\Omega^0(\mathfrak{g}_E)$ and set $K_0 = (1 + \nu_0[A_0]^{-1})(1 + \|F_{A_0}\|_{L^2})$. Let ε_1 be a constant satisfying $0 < \varepsilon_1 \leq zK_0^{-2}(1 + \nu_0[A_0]^{-1/2})^{-1}$. Then the following hold:*

1. *For any $[A] \in \mathcal{B}_E^k$ with $\text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) < \varepsilon_1$, there is a gauge transformation $u \in \mathcal{G}_E^{k+1}$ such that*
 - (a) $d_{A_0}^*(u(A) - A_0) = 0$,
 - (b) $\|u(A) - A_0\|_{L^{2\sharp,4}} \leq cK_0 \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0])$.
2. *For any $[A] \in \mathcal{B}_E^k$ with $\text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) < \varepsilon_1$, there is a gauge transformation $u \in \mathcal{G}_E^{k+1}$ such that*
 - (a) $d_{A_0}^*(u(A) - A_0) = 0$,
 - (b) $\|u(A) - A_0\|_{L^{2\sharp,4}} \leq cK_0 \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0])$,
 - (c) $\|u(A) - A_0\|_{L_{1,A_0}^2} \leq cK_0 \text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0])$.

Our first proof of Theorem 6.1, via the method of continuity, occupies Sections 6.1, 6.2 and 6.3. A rather different proof, via a direct application of the inverse function theorem using $L_2^{\sharp,2}$ gauge transformations, is given in Section 8.

6.1. Distance functions on the quotient space. Our first task is to verify the existence of minimizing gauge transformations $u \in \mathcal{G}_E^{k+1}$ for the family of distance functions on \mathcal{B}_E^k defined above: this is established in Lemma 6.3 and the proof uses the following version of Uhlenbeck's weak compactness theorem.

Proposition 6.2. *Let X be a closed, smooth, Riemannian four-manifold, let G be a compact Lie group, let M be a positive constant, let A_0 be an L_2^2 connection on a G bundle E over X . If $\{A_\alpha\}$ is a sequence of L_2^2 connections on E such that $\|F_{A_\alpha}\|_{L_{1,A_0}^2} \leq M$, then there is a subsequence $\{\alpha'\} \subset \{\alpha\}$ and a sequence of L_3^2 gauge transformations $\{u_{\alpha'}\}$ such that $u_{\alpha'}(A_{\alpha'})$ converges weakly in L_{2,A_0}^2 and strongly in L_{1,A_0}^p , for $1 \leq p < 4$, to an L_2^2 connection A_∞ on E .*

Proof. From the Sobolev embedding $L_1^2 \subset L^p$, $2 < p < 4$, we obtain a uniform L^p bound $\|F_{A_\alpha}\|_{L^p} \leq cM$ and so, according to [54, Theorem 3.6], there is a subsequence $\{\alpha'\} \subset \{\alpha\}$ and a sequence of L_2^p gauge transformations $\{u_{\alpha'}\}$ such that $u_{\alpha'}(A_{\alpha'})$ converges weakly in L_{1,A_0}^p to an L_1^p connection A_∞ on E . The stronger conclusion above is obtained simply

by reworking the proof of Theorem 3.6 in [54], using the following local estimate for the connections A_α over small balls $B \subset X$. Theorem 2.1 of [54] provides a sequence of local trivializations $v_\alpha : P|_B \rightarrow B \times G$ such that $a_\alpha = v_\alpha(A_\alpha) - \Gamma$ satisfies $d^*a_\alpha = 0$ and

$$\|a_\alpha\|_{L_1^p(B)} \leq c\|F_{A_\alpha}\|_{L^p(B)}, \quad 2 \leq p < 4,$$

where Γ is the product connection. Now $F_{A_\alpha} = da_\alpha + a_\alpha \wedge a_\alpha$, so

$$\begin{aligned} \|a_\alpha\|_{L_2^2(B)} &\leq \|da_\alpha\|_{L_1^2(B)} + \|a_\alpha\|_{L_1^2(B)} \\ &\leq \|a_\alpha \wedge a_\alpha\|_{L_1^2(B)} + \|F_{A_\alpha}\|_{L_1^2(B)} + \|a_\alpha\|_{L_1^2(B)}. \end{aligned}$$

Now, using the multiplication $L^6 \times L^3 \rightarrow L^2$, the embeddings $L_1^3 \subset L_1^{12/5} \subset L^6$ and $d(a_\alpha \wedge a_\alpha) = da_\alpha \wedge a_\alpha - a_\alpha \wedge da_\alpha$, we have

$$\|d(a_\alpha \wedge a_\alpha)\|_{L^2} \leq c\|da_\alpha\|_{L^3}\|a_\alpha\|_{L^6} \leq c\|a_\alpha\|_{L_1^3}^2,$$

while $\|a_\alpha \wedge a_\alpha\|_{L^2} \leq \|a_\alpha\|_{L^4(B)}^2 \leq c\|a_\alpha\|_{L_1^2(B)}^2$. Hence, we obtain

$$\begin{aligned} \|a_\alpha\|_{L_2^2(B)} &\leq c\|F_{A_\alpha}\|_{L_1^2(B)} + c\|a_\alpha\|_{L_1^3(B)}^2 + \|a_\alpha\|_{L_1^2(B)} \\ &\leq c\|F_{A_\alpha}\|_{L_1^2(B)}(1 + \|F_{A_\alpha}\|_{L_1^2(B)}) \\ &\leq c\|F_{A_\alpha}\|_{L_{1,A_0}^2(B)}(1 + \|F_{A_\alpha}\|_{L_{1,A_0}^2(B)})(1 + \|A_0 - \Gamma\|_{L_1^2}). \end{aligned}$$

In particular, the sequence of Coulomb-gauge, local connection matrices $\{a_\alpha\}$ is bounded in $L_2^2(B)$, so we can extract a weakly $L_2^2(B)$ -convergent and strongly $L_1^p(B)$ -convergent subsequence, via the compactness of embedding $L_2^2(B) \subset L_1^p(B)$ when $1 \leq p < 4$. The patching argument used to complete the proof of Uhlenbeck's theorem now proceeds exactly as in [54] to give the desired conclusion. \square

The proposition is used to extract the desired convergence in the next lemma.

Lemma 6.3. *For any points $[A_0], [A]$ in \mathcal{B}_E^k there are gauge transformations such that the following equalities hold:*

- (1) $\text{dist}_{L^4}([A], [A_0]) = \|u(A) - A_0\|_{L^4(X)}, \quad u \in \mathcal{G}_E^3$
- (2) $\text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) = \|v(A) - A_0\|_{\mathcal{L}_{1,A_0}^{\sharp,2}}, \quad v \in \mathcal{G}_E^3,$
- (3) $\text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) = \|w(A) - A_0\|_{L_{1,A_0}^{\sharp,2}}, \quad w \in \mathcal{G}_E^3,$
- (4) $\text{dist}_{L_{\ell,A_0}^2}([A], [A_0]) = \|w_\ell(A) - A_0\|_{L_{\ell,A_0}^2}, \quad \ell = 1 \text{ and } 3 \leq \ell \leq k,$

where $w_1 \in \mathcal{G}_E^3$ and $w_\ell \in \mathcal{G}_E^{\ell+1}$ in (4). Moreover, the above distance functions (including the $\ell = 2$ distance function in (4)) are continuous with respect to the quotient L_k^2 topology on \mathcal{B}_E^k .

Proof. Consider (1). Let $\{u_\alpha\}$ be a minimizing sequence in \mathcal{G}_E^{k+1} , so $\|u_\alpha(A) - A_0\|_{L^4}$ converges to $\text{dist}_{L^4}([A], [A_0])$ as $\alpha \rightarrow \infty$. Setting $B_\alpha = u_\alpha(A) = A - (d_A u_\alpha)u_\alpha^{-1} \in \mathcal{A}_E^k$, we see that $B_\alpha u_\alpha = Au_\alpha - d_A u_\alpha = Au_\alpha - d_{A_0} u_\alpha - [A - A_0, u_\alpha]$, and thus

$$(6.1) \quad d_{A_0} u_\alpha = u_\alpha(A - A_0) - (B_\alpha - A_0)u_\alpha.$$

Therefore, as $\|u_\alpha\|_{C^0} \leq c(G)$, we have

$$\|\nabla_{A_0} u_\alpha\|_{L^4} \leq c(\|A - A_0\|_{L^4} + \|B_\alpha - A_0\|_{L^4}),$$

so the sequence $\{u_\alpha\} \subset L^2_{k+1}(\mathfrak{gl}(E))$ is bounded in $L^4_{1,A_0}(\mathfrak{gl}(E))$. So, passing to a subsequence, we may suppose that $\{u_\alpha\}$ converges weakly in $L^4_{1,A_0}(\mathfrak{gl}(E))$ and strongly in $L^q(\mathfrak{gl}(E))$, via the compact embedding $L^4_1 \subset L^q$, for any $1 \leq q < \infty$, to a limit $u \in L^4_1(\mathfrak{gl}(E))$.

We also have $F_{B_\alpha} = F_{u_\alpha(A)} = u_\alpha F_A u_\alpha^{-1}$, so $\|F_{B_\alpha}\|_{L^2} = \|F_A\|_{L^2}$ and as

$$\nabla_{A_0} F_{B_\alpha} = (\nabla_{A_0} u_\alpha) \otimes F_A u_\alpha^{-1} + u_\alpha (\nabla_{A_0} F_A) u_\alpha^{-1} - u_\alpha F_A \otimes u_\alpha^{-1} (\nabla_{A_0} u_\alpha) u_\alpha^{-1},$$

we see that

$$(6.2) \quad \begin{aligned} \|\nabla_{A_0} F_{B_\alpha}\|_{L^2} &\leq c(\|\nabla_{A_0} u_\alpha\|_{L^4} \|F_A\|_{L^4} + \|\nabla_{A_0} F_A\|_{L^2}) \\ &\leq c(1 + \|u_\alpha\|_{L^4_{1,A_0}}) \|F_A\|_{L^2_{1,A_0}}. \end{aligned}$$

Hence, the sequence of L^2_k connections $\{B_\alpha\}$ has curvature uniformly bounded in L^2_{1,A_0} : Proposition 6.2 implies, after passing to a subsequence, that the sequence $\{B_\alpha\}$ converges weakly in L^2_{2,A_0} and strongly in L^p_{1,A_0} , for $1 \leq p < 4$, to an L^2_2 G connection B on E . From (6.1) we obtain

$$(6.3) \quad d_{A_0} u = u(A - A_0) - (B - A_0)u,$$

a first-order linear elliptic equation in u with L^2_2 coefficients. Therefore, $u \in L^2_3(\mathfrak{gl}(E))$ and $B = u(A) = A - (d_A u)u^{-1}$ lies in \mathcal{A}^2_E . It is not *a priori* clear that the limit u actually lies in \mathcal{G}^3_E (since the convergence was only weakly $L^4_{1,A_0}(\mathfrak{gl}(E))$ and strongly $L^q(\mathfrak{gl}(E))$): however, the argument of the last paragraph in the proof of Lemma 4.2.4 in [14, p. 130] applies (using the compactness of the structure group G) and shows that the limit gauge transformation u lies in \mathcal{G}^3_E . Since $B_\alpha = u_\alpha(A)$ converges strongly in L^p_{1,A_0} to $u(A)$ we now have

$$\text{dist}_{L^4}([A], [A_0]) = \lim_{\alpha \rightarrow \infty} \|u_\alpha(A) - A_0\|_{L^4} = \|u(A) - A_0\|_{L^4},$$

as required in (1). The same argument proves Assertions (2) and (3) and Assertion (4) when $\ell = 1$. The case $\ell \geq 3$ in (4) is straightforward as we can now apply Lemma 3.1 to obtain the desired convergence.

It remains to check L^2_k continuity. We just consider (1), as the remaining cases are identical. If $[A_\alpha] \in \mathcal{B}^k_E$ is a sequence converging to $[A_\infty] \in \mathcal{B}^k_E$, then there is a sequence of gauge transformations $s_\alpha \in \mathcal{G}^{k+1}_E$ such that $s_\alpha(A_\alpha)$ converges in L^2_{k,A_0} to $A_\infty \in \mathcal{A}^k_E$ and, in particular, in L^4 . But then

$$\begin{aligned} &|\text{dist}_{L^4}([A_\alpha], [A_0]) - \text{dist}_{L^4}([A_\infty], [A_0])| \\ &= |\text{dist}_{L^4}([s_\alpha(A_\alpha)], [A_0]) - \text{dist}_{L^4}([A_\infty], [A_0])| \\ &\leq \text{dist}_{L^4}([s_\alpha(A_\alpha)], [A_\infty]) \leq \|s_\alpha(A_\alpha) - A_\infty\|_{L^4}, \end{aligned}$$

and so

$$\lim_{\alpha \rightarrow \infty} \text{dist}_{L^4}([A_\alpha], [A_0]) = \text{dist}_{L^4}([A_\infty], [A_0]),$$

as desired. \square

6.2. Closedness. Let $\mathfrak{B} \subset \bar{B}^{1,\sharp,2}_{[A_0]}(\varepsilon)$ be the subset of points $[A]$ such that there exists a gauge transformation $u \in \mathcal{G}^{k+1}_E$ satisfying the conclusions of Assertion (2) of Theorem 6.1; let $\mathfrak{B}^* \subset \bar{B}^{1,*,2}_{[A_0]}(\varepsilon)$ be the subset of points $[A]$ such that there exists a gauge transformation $u \in \mathcal{G}^{k+1}_E$ satisfying the conclusions of Assertion (1). As in the proof of Theorem 2.1 in [54],

we apply the method of continuity to show that $\mathfrak{B}^* = \bar{B}_{[A_0]}^{1,*,2}(\varepsilon)$ and $\mathfrak{B} = \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon)$ for small enough ε . Not surprisingly, we have:

Lemma 6.4. *The balls $\bar{B}_{[A_0]}^{1,*,2}(\varepsilon)$ and $\bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon)$ are connected.*

Proof. If $[A] \in \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon)$, there is a gauge transformation $u \in \mathcal{G}_E^{k+1}$ such that $\|u(A) - A_0\|_{L_{1,A_0}^{\sharp,2}} \leq \varepsilon$. Then $A_t = A_0 + t(u(A) - A_0)$, $t \in [0, 1]$, is a path in \mathcal{A}_E^k joining A_0 to $u(A)$ and $\|A_t - A_0\|_{L_{1,A_0}^{\sharp,2}} = t\|u(A) - A_0\|_{L_{1,A_0}^{\sharp,2}} \leq t\varepsilon$, so the path $[A_t]$ lies in $\bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon)$ and joins $[A_0]$ to $[A]$. Similarly for $\bar{B}_{[A_0]}^{1,*,2}(\varepsilon)$. \square

Our task then reduces to showing that \mathfrak{B}^* is an open and closed subspace of $\bar{B}_{[A_0]}^{1,*,2}(\varepsilon)$ and that \mathfrak{B} is an open and closed subspace of $\bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon)$. First we consider closedness:

Lemma 6.5. *The subspaces $\mathfrak{B}^* \subset \bar{B}_{[A_0]}^{1,*,2}(\varepsilon)$ and $\mathfrak{B} \subset \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon)$ are closed.*

Proof. It suffices to consider the second assertion as the same argument yields the first. Suppose $[A_\alpha]$ is a sequence of points in \mathfrak{B} which converges in \mathcal{B}_E^k to a point $[B_\infty]$. We may suppose, without loss of generality, that $A_\alpha \in \mathcal{A}_E^k$ is the corresponding sequence of connections, representing the gauge-equivalence classes $[A_\alpha]$, which satisfy the defining conditions for \mathfrak{B} :

$$(6.4) \quad \begin{aligned} d_{A_0}^*(A_\alpha - A_0) &= 0, \\ \|A_\alpha - A_0\|_{L_{2,\sharp,4}^2} &\leq cK_0 \operatorname{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A_\alpha], [A_0]), \\ \|A_\alpha - A_0\|_{L_{1,A_0}^2} &\leq cK_0 \operatorname{dist}_{L_{1,A_0}^{\sharp,2}}([A_\alpha], [A_0]). \end{aligned}$$

Since $[A_\alpha]$ converges in \mathcal{B}_E^k to $[B_\infty]$, there is a sequence of gauge transformations $u_\alpha \in \mathcal{G}_E^{k+1}$ such that $B_\alpha := u_\alpha(A_\alpha)$ converges in L_{k,A_0}^2 to $B_\infty \in \mathcal{A}_E^k$. Since $B_\alpha = u_\alpha(A_\alpha)$ and $d_{A_0}^*(A_\alpha - A_0) = 0$, we have

$$(6.5) \quad d_{A_0} u_\alpha = u_\alpha(A_\alpha - A_0) - (B_\alpha - A_0)u_\alpha,$$

$$(6.6) \quad \begin{aligned} d_{A_0}^* d_{A_0} u_\alpha &= - * (d_{A_0} u_\alpha \wedge *(A_\alpha - A_0)) - (d_{A_0}^*(B_\alpha - A_0))u_\alpha \\ &\quad - * (* (B_\alpha - A_0) \wedge d_{A_0} u_\alpha), \end{aligned}$$

and so, as $\|u_\alpha\|_{C^0} \leq 1$,

$$\begin{aligned} \|d_{A_0} u_\alpha\|_{L_{2,\sharp,4}^2} &\leq \|A_\alpha - A_0\|_{L_{2,\sharp,4}^2} + \|B_\alpha - A_0\|_{L_{2,\sharp,4}^2} \\ \|d_{A_0}^* d_{A_0} u_\alpha\|_{L_{\sharp,2}^2} &\leq \|d_{A_0} u_\alpha\|_{L_{2,\sharp,4}^2} \|A_\alpha - A_0\|_{L_{2,\sharp,4}^2} + \|d_{A_0}^* B_\alpha - A_0\|_{L_{\sharp,2}^2} \\ &\quad \|B_\alpha - A_0\|_{L_{2,\sharp,4}^2} \|d_{A_0} u_\alpha\|_{L_{2,\sharp,4}^2}. \end{aligned}$$

Therefore, the sequence u_α is bounded in $L_{2,A_0}^2(\mathfrak{gl}(E))$ and so, passing to a subsequence, we may suppose that u_α converges weakly in $L_{2,A_0}^2(\mathfrak{gl}(E))$ (and strongly in L_{1,A_0}^p , for any $p < 4$ via the compact embedding $L_2^2 \subset L_1^p$) to a limit $u_\infty \in L^\infty \cap L_{2,A_0}^2(\mathfrak{gl}(E))$.

On the other hand, using $A_\alpha = u_\alpha^{-1}(B_\alpha)$, we have $\|F_{A_\alpha}\|_{L^2} = \|F_{B_\alpha}\|_{L^2}$ and the derivation of (6.2) gives

$$\|\nabla_{A_0} F_{A_\alpha}\|_{L^2} \leq c(1 + \|u_\alpha\|_{L_{1,A_0}^4}) \|F_{B_\alpha}\|_{L_{1,A_0}^2},$$

so the sequence A_α has curvature uniformly bounded in L^2_{1,A_0} . Thus, after passing to a subsequence we may assume by Proposition 6.2 that the sequence A_α converges weakly in L^2_{2,A_0} and strongly in L^p_{1,A_0} , $2 \leq p < 4$, to a limit $A_\infty \in \mathcal{A}^2_E$.

Taking weak limits in (6.5) and (6.6) yields

$$(6.7) \quad d_{A_0} u_\infty = u_\infty (A_\infty - A_0) - (B_\infty - A_0) u_\infty.$$

The equation (6.7) is first order, linear, elliptic in $u_\infty \in L^\infty \cap L^2_2$ with L^2_2 coefficients. Hence, u_∞ is in $L^2_3(\mathfrak{gl}(E))$ and in particular, in \mathcal{G}^3_E , while $B_\infty = u_\infty(A_\infty)$. From (6.7) we see that

$$A_\infty - A_0 = u_\infty^{-1} (B_\infty - A_0) u_\infty + u_\infty^{-1} d_{A_0} u_\infty$$

and so, as $d_{A_0}^*(A_\infty - A_0) = 0$, we have

$$(6.8) \quad d_{A_0}^*(u_\infty^{-1} d_{A_0} u_\infty + u_\infty^{-1} (B_\infty - A_0) u_\infty) = 0.$$

This is a second-order elliptic equation for $u_\infty \in \mathcal{G}^3_E$ with L^2_k coefficients: in particular, since $u_\infty \in L^p_2$ for $2 \leq p \leq 4$, a standard elliptic bootstrapping argument then implies that $u_\infty \in L^2_{k+1}$ (see, for example, the proof of Proposition 3.3 in [16]) and therefore $A_\infty = u_\infty^{-1} (B_\infty) \in \mathcal{A}^k_E$.

Now, taking weak limits in (6.4), we have

$$\begin{aligned} d_{A_0}^*(A_\infty - A_0) &= \lim_{\alpha \rightarrow \infty} d_{A_0}^*(A_\alpha - A_0) = 0, \\ \|A_\infty - A_0\|_{L^{2\sharp,4}} &= \lim_{\alpha \rightarrow \infty} \|A_\alpha - A_0\|_{L^{2\sharp,4}} \leq \lim_{\alpha \rightarrow \infty} cK_0 \operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A_\alpha], [A_0]), \\ \|A_\alpha - A_0\|_{L^{\sharp,2}_{1,A_0}} &= \lim_{\alpha \rightarrow \infty} \|A_\alpha - A_0\|_{L^{\sharp,2}_{1,A_0}} \leq \lim_{\alpha \rightarrow \infty} cK_0 \operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A_\alpha], [A_0]). \end{aligned}$$

Moreover, as $B_\infty = u_\infty(A_\infty)$ and $u_\infty \in \mathcal{G}^{k+1}_E$,

$$\begin{aligned} \lim_{\alpha \rightarrow \infty} \operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A_\alpha], [A_0]) &= \operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([B_\infty], [A_0]) = \operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A_\infty], [A_0]), \\ \lim_{\alpha \rightarrow \infty} \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A_\alpha], [A_0]) &= \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([B_\infty], [A_0]) = cK_0 \operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A_\infty], [A_0]), \end{aligned}$$

where the L^2_k continuity of the distance functions is given by Lemma 6.3. Therefore, $[B_\infty] = [A_\infty] \in \mathfrak{B}$. Thus, \mathfrak{B} is closed in \mathcal{B}^k_E and in particular, closed in $\bar{B}^{1,\sharp,2}_{[A_0]}(\varepsilon)$, as desired. \square

6.3. Openness. We must first compare distances from the connection A_0 in the Coulomb slice through A_0 in \mathcal{A}^k_E and gauge-invariant distances in \mathcal{B}^k_E from the point $[A_0]$:

Lemma 6.6. *Let (X, g) be a closed, smooth, Riemannian four-manifold. Then there are positive constants c, z with the following significance. Let $K_0 = (1 + \nu_0[A_0]^{-1})(1 + \|F_{A_0}\|_{L^2})$. If $A \in \mathcal{A}^k_E$ satisfies*

- $d_{A_0}^*(A - A_0) = 0$,
- $\|A - A_0\|_{L^{2\sharp,4}} \leq zK_0^{-1}$,

then the following hold:

1. *If $\operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A], [A_0]) \leq zK_0^{-1}$, then*

$$\|A - A_0\|_{L^{2\sharp,4}} \leq cK_0 \operatorname{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A], [A_0]);$$

2. If $\text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) \leq zK_0^{-1}$, then

$$\begin{aligned} \|A - A_0\|_{L^{2\sharp,4}} &\leq cK_0 \text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]), \\ \|A - A_0\|_{L_{1,A_0}^2} &\leq cK_0 \text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]). \end{aligned}$$

Proof. Recall that for either distance function, minimizing gauge transformations in \mathcal{G}_E^3 exist by Lemma 6.3; for convenience, we denote both by $u \in \mathcal{G}_E^3$ although they need not *a priori* coincide. Since $B := u(A) = A - (d_A u)u^{-1} \in \mathcal{A}_E^2$, we have

$$u(A) - A_0 = u(A - A_0)u^{-1} - (d_{A_0} u)u^{-1}.$$

Our task, in essence, is to estimate the second term on the right above. Rewriting this equality gives a first-order, linear elliptic equation in u with L_2^2 coefficients:

$$(6.9) \quad d_{A_0} u = u(A - A_0) - (B - A_0)u.$$

Let $u_0 \in L_3^2(\mathfrak{gl}(E))$ be the L^2 orthogonal projection of $u \in \mathcal{G}_E^3 \subset L_3^2(\mathfrak{gl}(E))$ onto $\text{Ker}(d_{A_0}|_{L_3^2})^\perp$, so $u = u_0 + \gamma$, where $\gamma \in \text{Ker } d_{A_0} \subset \Omega^0(\mathfrak{gl}(E))$. Thus, as $d_{A_0}^*(A - A_0) = 0$ and $d_{A_0} u = d_{A_0} u_0$, we see that

$$\begin{aligned} d_{A_0}^* d_{A_0} u_0 &= - * (d_{A_0} u \wedge *(A - A_0)) + u d_{A_0}^*(A - A_0) \\ &\quad - (d_{A_0}^*(B - A_0))u - *(*(B - A_0) \wedge d_{A_0} u) \\ &= - * (d_{A_0} u_0 \wedge *(A - A_0)) - (d_{A_0}^*(B - A_0))u - *(*(B - A_0) \wedge d_{A_0} u_0). \end{aligned}$$

Therefore, using the bound $\|u\|_{C^0} \leq 1$ for any $u \in \mathcal{G}_E^3$ (as the representation for G is orthogonal), we have

$$\begin{aligned} \|\Delta_{A_0}^0 u_0\|_{L^{\sharp,2}} &\leq \|d_{A_0} u_0\|_{L^{2\sharp,4}} \|A - A_0\|_{L^{2\sharp,4}} + \|d_{A_0}^*(B - A_0)\|_{L^{\sharp,2}} \|u\|_{C^0} \\ &\quad + \|B - A_0\|_{L^{2\sharp,4}} \|d_{A_0} u_0\|_{L^{2\sharp,4}} \\ &\leq C (\|A - A_0\|_{L^{2\sharp,4}} + \|B - A_0\|_{L^{2\sharp,4}}) \|d_{A_0}^* d_{A_0} u_0\|_{L^{\sharp,2}} \\ &\quad + \|d_{A_0}^*(B - A_0)\|_{L^{\sharp,2}}, \end{aligned}$$

where $C = cK_0$. Now $\|B - A_0\|_{L^{2\sharp,4}} \leq c\|B - A_0\|_{L_{1,A_0}^2}$ via the embedding $L_1^2 \subset L^{2\sharp,4}$ of Lemma 4.1. For either $\text{dist}_{\mathcal{L}_{1,A_0}^{\sharp,2}}([A], [A_0]) \leq \frac{1}{4}C^{-1}$ or $\text{dist}_{L_{1,A_0}^{\sharp,2}}([A], [A_0]) \leq \frac{1}{4}C^{-1}$ and $\|A - A_0\|_{L^{2\sharp,4}} \leq \frac{1}{4}C^{-1}$, rearrangement yields

$$(6.10) \quad \|\Delta_{A_0}^0 u_0\|_{L^{\sharp,2}} \leq 2\|d_{A_0}^*(B - A_0)\|_{L^{\sharp,2}}.$$

On the other hand, from Lemma 5.6 we have

$$(6.11) \quad \begin{aligned} \|u_0\|_{L_{2,A_0}^{\sharp,2}} &\leq C\|\Delta_{A_0}^0 u_0\|_{L^{\sharp,2}}, \\ \|u_0\|_{L_{1,A_0}^4} &\leq C\|\Delta_{A_0}^0 u_0\|_{L^{\sharp,2}}, \end{aligned}$$

where $C = cK_0$ and the second bound follows from the embedding $L_2^2 \subset L_1^4$. So, combining (6.10) and (6.11) yields:

$$(6.12) \quad \begin{aligned} \|u_0\|_{L_{2,A_0}^{\sharp,2}} &\leq C\|d_{A_0}^*(B - A_0)\|_{L^{\sharp,2}}, \\ \|u_0\|_{L_{1,A_0}^4} &\leq C\|d_{A_0}^*(B - A_0)\|_{L^{\sharp,2}}. \end{aligned}$$

Consequently, using $d_{A_0}u = d_{A_0}u_0$ and (6.9) rewritten in the form,

$$(6.13) \quad u^{-1}(B - A_0)u = (A - A_0) - u^{-1}d_{A_0}u_0,$$

we obtain

$$(6.14) \quad \|A - A_0\|_{L^{2\sharp,4}} \leq \|u^{-1}(B - A_0)u\|_{L^{2\sharp,4}} + \|u^{-1}d_{A_0}u_0\|_{L^{2\sharp,4}},$$

$$(6.15) \quad \|A - A_0\|_{L^2_{1,A_0}} \leq \|u^{-1}(B - A_0)u\|_{L^2_{1,A_0}} + \|u^{-1}d_{A_0}u_0\|_{L^2_{1,A_0}}.$$

From (6.14) and (6.12), we see that

$$\begin{aligned} \|A - A_0\|_{L^{2\sharp,4}} &\leq \|B - A_0\|_{L^{2\sharp,4}} + \|d_{A_0}u_0\|_{L^{2\sharp,4}} \\ &\leq \text{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A], [A_0]) + C\|d_{A_0}^*(B - A_0)\|_{L^{\sharp,2}} \\ &\leq (1 + C) \text{dist}_{\mathcal{L}^{\sharp,2}_{1,A_0}}([A], [A_0]), \end{aligned}$$

giving the desired $L^{2\sharp,4}$ estimate for $A - A_0$.

Considering the first term in (6.13), we have

$$\begin{aligned} \nabla_{A_0}(u^{-1}(B - A_0)u) &= -u^{-1}(\nabla_{A_0}u)u^{-1} \otimes (B - A_0)u + u^{-1}(\nabla_{A_0}(B - A_0))u \\ &\quad + u^{-1}(B - A_0) \otimes \nabla_{A_0}u, \end{aligned}$$

and so applying (6.12), noting that $\nabla_{A_0}u = \nabla_{A_0}u_0$ and $\|u\|_{C^0} \leq 1$, we have

$$\begin{aligned} \|\nabla_{A_0}(u^{-1}(B - A_0)u)\|_{L^2} &\leq \|\nabla_{A_0}u_0\|_{L^4}\|B - A_0\|_{L^4} + \|\nabla_{A_0}(B - A_0)\|_{L^2} \\ &\leq C \text{dist}_{L^{\sharp,2}_{1,A_0}}^2([A], [A_0]) + \text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]). \end{aligned}$$

Thus, if $\text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) \leq \frac{1}{4}C^{-1}$, say, we obtain

$$(6.16) \quad \|\nabla_{A_0}(u^{-1}(B - A_0)u)\|_{L^2} \leq 2 \text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]).$$

Similarly, considering the second term in (6.13), we have

$$\nabla_{A_0}(u^{-1}d_{A_0}u_0) = -u^{-1}(\nabla_{A_0}u)u^{-1} \otimes d_{A_0}u_0 + u^{-1}\nabla_{A_0}d_{A_0}u_0$$

and therefore, by (6.12), we see that

$$\begin{aligned} \|\nabla_{A_0}(u^{-1}d_{A_0}u_0)\|_{L^2} &\leq \|\nabla_{A_0}u_0\|_{L^4}^2 + \|\nabla_{A_0}^2u_0\|_{L^2} \\ &\leq C \text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) \left(1 + C \text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0])\right). \end{aligned}$$

Provided $\text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) \leq \frac{1}{4}C^{-1}$, we obtain

$$(6.17) \quad \|\nabla_{A_0}(u^{-1}d_{A_0}u_0)\|_{L^2} \leq 2C \text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]).$$

Taking the L^2 norm of (6.13) and applying (6.12) to estimate the second term gives

$$\begin{aligned} \|A - A_0\|_{L^2} &\leq \|B - A_0\|_{L^2} + \|d_{A_0}u_0\|_{L^2} \\ &\leq \text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) + C\|d_{A_0}^*(B - A_0)\|_{L^{\sharp,2}}, \end{aligned}$$

and so

$$(6.18) \quad \|A - A_0\|_{L^2} \leq (1 + C) \text{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]).$$

Combining the estimates (6.15), (6.16), (6.17), and (6.18) yields

$$\|A - A_0\|_{L^2_{1,A_0}} \leq 2(1 + C) \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]),$$

giving the desired L^2_{1,A_0} estimate for $A - A_0$. \square

Naturally, a comparison — going in the other direction — of distances from A_0 in the Coulomb slice in \mathcal{A}_E^k through A_0 and gauge-invariant distances in \mathcal{B}_E^k from the point $[A_0]$ is elementary: If $A \in \mathbf{S}_{A_0}$ and $\|A - A_0\|_{L^2_{k,A_0}} < \delta$, say, then Lemma 4.1 implies that

$$(6.19) \quad \begin{aligned} \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) &\leq c\|A - A_0\|_{L^2_{k,A_0}} < c\delta, & k \geq 1, \\ \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) &\leq c\|A - A_0\|_{L^2_{k,A_0}} < c\delta, & k \geq 2, \end{aligned}$$

for some positive constant $c(X, g, k)$. The observation is used in concluding that \mathfrak{B}^* , \mathfrak{B} are open subspaces of $\bar{B}_{[A_0]}^{1,*}, \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon_1)$, respectively:

Lemma 6.7. *Let (X, g) be a closed, smooth, Riemannian four-manifold and let G be a compact Lie group. Then there is a positive constant z with the following significance. Let $K_0 = (1 + \nu_0[A_0]^{-1})(1 + \|F_{A_0}\|_{L^2})$. If $\varepsilon_1 < zK_0^{-2}(1 + \nu_0[A_0]^{-1/2})^{-1}$, then the following hold:*

- \mathfrak{B}^* is an open subspace of $\bar{B}_{[A_0]}^{1,*}(\varepsilon_1)$;
- \mathfrak{B} is an open subspace of $\bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon_1)$.

Proof. It suffices to consider the second assertion, as the argument for the first is identical. Suppose $[A] \in \mathfrak{B}$ and that $A \in \mathcal{A}_E^k$ is a representative satisfying the defining conditions for \mathfrak{B} . Then A satisfies $d_{A_0}^*(A - A_0) = 0$ and the estimates

$$\begin{aligned} \|A - A_0\|_{L^{2\sharp,4}} &\leq c_0 K_0 \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) \leq c_0 K_0 \varepsilon_1 \\ \|A - A_0\|_{L^2_{1,A_0}} &\leq c_0 K_0 \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) \leq c_0 K_0 \varepsilon_1, \end{aligned}$$

while $\|A - A_0\|_{L^{2\sharp,4}} \leq c_1 \|A - A_0\|_{L^2_{1,A_0}}$ via the Sobolev embedding $L^2_1 \subset L^{2\sharp,4}$ and Kato's inequality. Consequently, if $c_1 c_0 K_0 \varepsilon_1 \leq \frac{1}{2} \varepsilon_0$, then $A \in \mathbf{B}_{A_0}^4(\varepsilon_0) \subset \mathcal{A}_E^k$ and we see that

$$\bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon_1) \subset \pi(\mathbf{B}_{[A_0]}^4(\varepsilon_0)).$$

Lemma 3.6 implies that the map $\pi : \mathbf{B}_{[A_0]}^4(\varepsilon_0) / \operatorname{Stab}_{A_0} \rightarrow \mathcal{B}_E^k$ given by $A' \mapsto [A']$ is a local homeomorphism onto its image $\pi(\mathbf{B}_{[A_0]}^4(\varepsilon_0))$ for any $\varepsilon_0 < z(1 + \nu_0[A_0]^{-1/2})^{-1}$. In particular, if $A' \in \mathbf{B}_{[A_0]}^4(\varepsilon_0)$ and $\|A' - A\|_{L^2_{k,A_0}} < \delta$, then $A' \in B_{[A_0]}^{1,\sharp,2}(\varepsilon_1) \subset \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon_1)$ for small enough δ .

The embedding $L^2_1 \subset L^{2\sharp,4}$ and Lemma 6.6 imply that if $\|A' - A_0\|_{L^2_{1,A_0}} \leq zK_0^{-1}$ and $\operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A'], [A_0]) \leq zK_0^{-1}$, then

$$\begin{aligned} \|A' - A_0\|_{L^{2\sharp,4}} &\leq cK_0 \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A'], [A_0]) \leq cK_0 \varepsilon_1, \\ \|A' - A_0\|_{L^2_{1,A_0}} &\leq cK_0 \operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A'], [A_0]) \leq cK_0 \varepsilon_1. \end{aligned}$$

These inequalities are satisfied by A ; moreover $\operatorname{dist}_{L^{\sharp,2}_{1,A_0}}([A], [A_0]) \leq \varepsilon_1$ and $\|A - A_0\|_{L^2_{1,A_0}} \leq c_0 K \varepsilon_1$. Require that $\varepsilon_1 \leq \frac{1}{2} z K_0^{-1}$ and $c_0 K \varepsilon_1 \leq \frac{1}{2} z K_0^{-1}$, so $\varepsilon_1 \leq \frac{1}{2} z \min\{1, c_0\} K_0^{-2}$. Hence,

if A' is L_{k,A_0}^2 -close enough to A (where $k \geq 2$), we can ensure $[A']$ obeys the last two defining conditions for \mathfrak{B} and so $[A'] \in \mathfrak{B}$. Thus, $\mathfrak{B} \subset \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon_1)$ is open, as desired. \square

We can now conclude the proof of Theorem 6.1:

Proof of Theorem 6.1. Lemmas 6.5 and 6.7 imply that \mathfrak{B} is an open and closed subset of the connected space $\bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon_1)$, so $\mathfrak{B} = \bar{B}_{[A_0]}^{1,\sharp,2}(\varepsilon_1)$. Similarly for \mathfrak{B}^* and $\bar{B}_{[A_0]}^{1,*,2}(\varepsilon_1)$ and hence the result follows. \square

Similarly, we conclude the proof of Theorem 1.1:

Proof of Theorem 1.1. Given Theorem 6.1, the only assertion left unaccounted for is the uniqueness of the gauge transformation $u \in \mathcal{G}_E^{k+1}$, modulo Stab_{A_0} . But this follows from Lemma 3.7, just as in the paragraph immediately following the proof of that lemma. \square

7. CRITICAL-EXPONENT SOBOLEV NORMS AND THE GROUP OF GAUGE TRANSFORMATIONS

We now define an $L_2^{\sharp,2}$ space of gauge transformations, by analogy with the definition of \mathcal{G}_E^{k+1} when $k \geq 2$, and set

$$\mathcal{G}_E^{2,\sharp,2} := \{u \in L_2^{\sharp,2}(\mathfrak{gl}(E)) : u \in G \text{ a.e.}\} \subset L_k^2(\mathfrak{gl}(E)).$$

It is not entirely clear *a priori* that $\mathcal{G}_E^{2,\sharp,2}$ is a Banach Lie group. In the case of its counterpart, \mathcal{G}_E^{k+1} , the manifold structure follows from the fact that the exponential map

$$\text{Exp} : T_{\text{id}_E} \mathcal{G}_E = \Omega^0(\mathfrak{g}_E) \rightarrow \mathcal{G}_E, \quad \zeta \mapsto \text{Exp } \zeta,$$

extends to a smooth map $\text{Exp} : L_{k+1}^2(\mathfrak{g}_E) \rightarrow L_{k+1}^2(\mathfrak{g}_E)$ and defines a system of smooth coordinate charts for \mathcal{G}_E^{k+1} . Here, Exp is defined pointwise at $u \in \mathcal{G}_E$ for $\zeta \in T_{\text{id}_E} \mathcal{G}_E$ by setting

$$(\text{Exp}_u \zeta)(x) := \exp_{u(x)}(\zeta(x)), \quad x \in X,$$

where $\exp : \mathfrak{g} \rightarrow G$ is the usual, C^∞ exponential map for the Lie group G on the right-hand side [22, Appendix A].

To verify that $\mathcal{G}_E^{2,\sharp,2}$ is in fact a Banach Lie group we will need estimates for the covariant derivatives of the exponential map. The estimates below follow by reworking the usual proof of the Sobolev lemma for left composition of Sobolev sections by smooth vector bundle maps [37, Lemma 9.9]. The difference here is that we keep track of the dependence of the constants on the geometric data: this precision is required for the implicit function argument in the next section in order to complete the proof of our slice theorem.

For $\chi, \zeta, \xi \in \Omega^0(\mathfrak{g}_E)$, the differentials

$$\begin{aligned} (D \text{Exp})_\chi : \Omega^0(\mathfrak{g}_E) &\rightarrow T_{\text{Exp } \chi} \mathcal{G}_E, & \zeta &\mapsto (D \text{Exp})_\chi \zeta, \\ (D^2 \text{Exp})_{\chi, \zeta} : \Omega^0(\mathfrak{g}_E) &\rightarrow T_{\text{Exp } \chi} \mathcal{G}_E, & \xi &\mapsto (D^2 \text{Exp})_{\chi, \zeta} \xi, \end{aligned}$$

are defined pointwise by setting

$$\begin{aligned} (D \text{Exp})_\chi \zeta|_x &= (D \exp)_{\chi(x)} \zeta(x), \\ (D^2 \text{Exp})_{\chi, \zeta} \xi|_x &= (D^2 \exp)_{\chi(x), \zeta(x)} \xi(x), \end{aligned}$$

for any $x \in X$. When writing the differential $(D^2 \text{Exp})_{\chi, \zeta}$ above, we have identified $T_{(D \text{Exp})_\chi} (T_{\text{Exp } \chi} \mathcal{G}_E)$ with $T_{\text{Exp } \chi} \mathcal{G}_E$.

The maps $(D \operatorname{Exp})_\chi : \Omega^0(\mathfrak{g}_E) \rightarrow \Omega^0(\mathfrak{g}_E)$ and $(D^2 \operatorname{Exp})_{\chi, \zeta} : \Omega^0(\mathfrak{g}_E) \rightarrow \Omega^0(\mathfrak{g}_E)$ extend linearly to maps

$$\begin{aligned} (D \operatorname{Exp})_\chi &: C^\infty(\otimes^\ell(T^*X) \otimes \mathfrak{g}_E) \rightarrow C^\infty(\otimes^\ell(T^*X) \otimes \mathfrak{g}_E), \\ (D^2 \operatorname{Exp})_\chi &: C^\infty(\otimes^\ell(T^*X) \otimes \mathfrak{g}_E) \rightarrow C^\infty(\otimes^\ell(T^*X) \otimes \mathfrak{g}_E), \end{aligned}$$

for $\ell \geq 1$, by setting

$$\begin{aligned} (D \operatorname{Exp})_\chi(\theta \otimes \zeta) &= \theta \otimes (D \operatorname{Exp})_\chi \zeta, \\ (D^2 \operatorname{Exp})_{\chi, \zeta}(\theta \otimes \xi) &= \theta \otimes (D^2 \operatorname{Exp})_{\chi, \zeta} \xi, \end{aligned}$$

for $\theta \in \otimes^\ell(T^*X)$ and $\xi \in \Omega^0(\mathfrak{g}_E)$. As usual, we embed $\mathcal{G}_E \subset \Omega^0(\mathfrak{g}_E)$ in order to compute the covariant derivatives of sections $u \in \mathcal{G}_E$.

Lemma 7.1. *Let G be a compact Lie group. Then there is a positive constant $c(G)$ with the following significance. Let X be a closed, smooth, Riemannian four-manifold. If A is a C^∞ connection on a G bundle E , and $\chi \in \Omega^0(\mathfrak{g}_E)$, then $e^\chi \in \mathcal{G}_E$ satisfies the following pointwise bounds:*

$$\begin{aligned} (1) \quad & |\nabla_A e^\chi| \leq |\nabla_A \chi| + c|\chi||\nabla_A \chi|, \\ (2) \quad & |\nabla_A^2 e^\chi| \leq c(|\chi| + |\nabla_A \chi|)|\nabla_A \chi| + c(1 + |\chi|)|\nabla_A^2 \chi|, \\ (3) \quad & |\nabla_A^* \nabla_A e^\chi| \leq c(|\chi| + |\nabla_A \chi|)|\nabla_A \chi| + c(1 + |\chi|)|\nabla_A^* \nabla_A \chi|. \end{aligned}$$

Proof. We have

$$\nabla_A e^\chi = \nabla_A(\operatorname{Exp} \chi) = (D \operatorname{Exp})_\chi \circ d_A \chi \in \Omega^1(\mathfrak{g}_E).$$

Since $(D \operatorname{exp})_0 = \operatorname{id}_\mathbb{E}$ and $\operatorname{exp} : \mathfrak{g} \rightarrow G$ is analytic, we have the pointwise bound $|(D \operatorname{exp})_{\chi(x)} - \operatorname{id}_\mathbb{E}| \leq c(G)|\chi(x)|$ and thus a pointwise bound

$$|(D \operatorname{Exp})_\chi - \operatorname{id}_E| \leq c|\chi|,$$

noting that $(D \operatorname{Exp})_0 = \operatorname{id}_E$. Therefore, we have

$$|\nabla_A e^\chi| \leq |\nabla_A \chi| + c|\chi||\nabla_A \chi|,$$

which gives the first assertion.

Define $\Phi(\chi, \zeta) = (D \operatorname{Exp})_\chi(\zeta) \in \Omega^1(\mathfrak{g}_E)$, for $\chi \in \Omega^0(\mathfrak{g}_E)$ and $\zeta \in \Omega^1(\mathfrak{g}_E)$, noting that Φ is nonlinear in χ , but linear in ζ . Thus,

$$\nabla_A^2 u = (D_1 \Phi)_{(\chi, \nabla_A \chi)}(\nabla_A \chi) + (D_2 \Phi)_\chi(\nabla_A^2 \chi),$$

where $D_i \Phi$, $i = 1, 2$, denote the partial derivatives of Φ with respect to first and second variables. Since $(D \Phi)_{(0,0)} = (D^2 \operatorname{Exp})_{(0,0)} = \operatorname{id}_E$, as $(D^2 \operatorname{exp})_{0,0} = \operatorname{id}_\mathbb{E}$, and $\operatorname{exp} : \mathfrak{g} \rightarrow G$ is analytic we have the pointwise bound

$$|\nabla_A^2 u| \leq c(|\chi| + |\nabla_A \chi|)|\nabla_A \chi| + c(1 + |\chi|)|\nabla_A^2 \chi|,$$

giving the second assertion. Similarly, as $*\Phi(\chi, \zeta) = \Phi(\chi, *\zeta)$ and $\nabla_A^* \nabla_A u = -*\nabla_A * \nabla_A u$, we have

$$|\nabla_A^* \nabla_A u| \leq c(|\chi| + |\nabla_A \chi|)|\nabla_A \chi| + c(1 + |\chi|)|\nabla_A^* \nabla_A \chi|,$$

giving the third assertion. \square

The preceding pointwise bounds for $\nabla_A u$, $\nabla_A^2 u$, and $\nabla_A^* \nabla_A u$ yield the following estimates for the exponential map:

Lemma 7.2. *Let G be a compact Lie group. Then there is a positive constant $c(G)$ with the following significance. Let X be a closed, smooth, Riemannian four-manifold. If $k \geq 2$ is an integer (so $L_{k+1}^2 \subset C^0$), A is an L_k^2 connection on a G bundle E , and $\chi \in L_{k+1}^2(\mathfrak{g}_E)$, then $e^\chi \in \mathcal{G}_E^{k+1}$ satisfies*

$$\begin{aligned}
(1) \quad & \|\nabla_A e^\chi\|_{L^2(X)} \leq \|\nabla_A \chi\|_{L^2(X)} + c\|\chi\|_{C^0(X)}\|\nabla_A \chi\|_{L^2(X)}, \\
(2) \quad & \|\nabla_A e^\chi\|_{L^\sharp(X)} \leq \|\nabla_A \chi\|_{L^\sharp(X)} + c\|\chi\|_{C^0(X)}\|\nabla_A \chi\|_{L^\sharp(X)}, \\
(3) \quad & \|\nabla_A^2 e^\chi\|_{L^2(X)} \leq c\|\chi\|_{C^0(X)}\|\nabla_A \chi\|_{L^2(X)} + \|\nabla_A \chi\|_{L^4(X)}^2 \\
& \quad + c(1 + \|\chi\|_{C^0(X)})\|\nabla_A^2 \chi\|_{L^2(X)}, \\
(4) \quad & \|\nabla_A^* \nabla_A e^\chi\|_{L^\sharp(X)} \leq c\|\chi\|_{C^0(X)}\|\nabla_A \chi\|_{L^\sharp(X)} + \|\nabla_A \chi\|_{L^{2^\sharp}(X)}^2 \\
& \quad + c(1 + \|\chi\|_{C^0(X)})\|\nabla_A^* \nabla_A \chi\|_{L^\sharp(X)},
\end{aligned}$$

Moreover, the preceding bounds continue to hold for $\chi \in L_2^{\sharp,2}(\mathfrak{g}_E) \subset C^0(\mathfrak{g}_E)$, with A an $L_1^{\sharp,2}$ connection on E , and $\text{Exp} : \Omega^0(\mathfrak{g}_E) \rightarrow \mathcal{G}_E$ extends to a continuous map $\text{Exp} : L_2^{\sharp,2}(\mathfrak{g}_E) \rightarrow \mathcal{G}_E^{2,\sharp,2}$.

Let $\mathcal{A}_E^{1,\sharp,2} = A_0 + L_{1,A_0}^{\sharp,2}(\Lambda^1 \otimes \mathfrak{g}_E)$, for any C^∞ reference connection A_0 on E . Recall that we have an embedding $L_2^{\sharp,2}(\mathfrak{g}_E) \subset C^0(\mathfrak{g}_E)$ and that the space $L_2^{\sharp,2}(\mathfrak{g}_E)$ is an algebra, while $L_1^{\sharp,2}(\Lambda^1 \otimes \mathfrak{g}_E)$ and $L_1^2(\Lambda^1 \otimes \mathfrak{g}_E)$ are $L_2^{\sharp,2}(\mathfrak{g}_E)$ -modules. Therefore, the proofs of Propositions (A.2) and (A.3) in [22] extend easily to give the following analogue for $\mathcal{G}_E^{2,\sharp,2}$ in place of \mathcal{G}_E^{k+1} :

Lemma 7.3. *Let X be a closed Riemannian four-manifold and let E be a Hermitian vector bundle over X . Then the following hold.*

1. *The space $\mathcal{G}_E^{2,\sharp,2}$ is a Banach Lie group with Lie algebra $T_{\text{id}_E} \mathcal{G}_E^{2,\sharp,2} = L_2^{\sharp,2}(\mathfrak{g}_E)$;*
2. *The action of $\mathcal{G}_E^{2,\sharp,2}$ on \mathcal{A}_E^1 and $\mathcal{A}_E^{1,\sharp,2}$ is smooth;*
3. *For $A \in \mathcal{A}_E^{1,\sharp,2}$, the differential, at the identity $\text{id}_E \in \mathcal{G}_E^{2,\sharp,2}$, of the map $\mathcal{G}_E^{2,\sharp,2} \rightarrow \mathcal{A}_E^{1,\sharp,2}$ given by $u \mapsto u(A) = A - (d_A u)u^{-1}$ is $\zeta \mapsto -d_A \zeta$ as a map $L_2^{\sharp,2}(\mathfrak{g}_E) \rightarrow L_1^{\sharp,2}(\Lambda^1 \otimes \mathfrak{g}_E)$, and similarly for $A \in \mathcal{A}_E^1$.*

8. EXISTENCE OF GAUGE TRANSFORMATIONS VIA THE INVERSE FUNCTION THEOREM

Our goal in this section is to give an alternative, ‘direct’ proof of Theorem 6.1 via the inverse function theorem. A direct argument — due to our overarching constraint of ensuring that the constants given there ultimately depend at most on the L^2 norm of the curvature and the least positive eigenvalue $\nu_0[A_0]$ — appears to be difficult within the standard framework of L_2^p ($p > 2$) gauge transformations acting on L_1^p connections; of course, if this constraint is dropped then a direct proof is standard. On the other hand, with the results of the last section, it is fairly straightforward within the framework of $L_2^{\sharp,2}$ gauge transformations.

We already know that $\pi(\mathbf{B}_{A_0}^4(\varepsilon_0))$ is open in \mathcal{B}_E^k , so it necessarily contains an L_{k,A_0}^2 -ball centered at $[A_0]$. Via the inverse function theorem we estimate the radii of $\mathcal{L}_{1,A_0}^{\sharp,2}$ and $L_{1,A_0}^{\sharp,2}$ balls, $B_{[A_0]}^{1,\sharp,2}(\varepsilon)$ and $B_{[A_0]}^{1,\sharp,2}(\varepsilon)$, which are contained in $\pi(\mathbf{B}_{A_0}^4(\varepsilon_0))$. Let us first dispose of the question of regularity for solutions to the second-order gauge-fixing equation:

Lemma 8.1. *Let X be a closed, Riemannian four-manifold. Then there is a constant ε with the following significance. Let G be compact Lie group and let $k \geq 2$ be an integer. Suppose that A_0 is an L_k^2 connection on a G bundle E , that $a \in L_k^2(\Lambda^1 \otimes \mathfrak{g}_E)$ and $\chi \in L_2^\sharp(\mathfrak{g}_E)$, and that $u = e^\chi$ is a solution to*

$$d_{A_0}^* ((d_{A_0} u) u^{-1} - u a u^{-1}) = 0.$$

If $\|d_{A_0} u\|_{L^4} < \varepsilon$ then $\chi \in L_{k+1}^2(\mathfrak{g}_E)$ and $u = e^\chi \in \mathcal{G}_E^{k+1}$.

Proof. Differentiation and right multiplication by u yields

$$(8.1) \quad d_{A_0}^* d_{A_0} u + *((d_{A_0} u) \wedge u^{-1} d_{A_0} u) + *(d_{A_0} u \wedge *a) + u d_{A_0}^* a + *(u a \wedge *u^{-1} d_{A_0} u) = 0.$$

From Lemma 4.2 we know that $u \in C^0 \cap L_1^2$ and so the last four terms in (8.1) are in L^2 . Hence, $d_{A_0}^* d_{A_0} u$ is in L^2 and so $u \in L_2^2$ by elliptic regularity for $d_{A_0}^* d_{A_0}$. The Sobolev embedding $L_1^2 \subset L^4$ and multiplication $L^4 \times L^q \rightarrow L^p$ for $2 \leq p < 4$ and $1/p = 1/4 + 1/q$ (so $4 \leq q < \infty$) now show that the last three terms in (8.1) are in L^p , so the equation takes the simpler form

$$(8.2) \quad d_{A_0}^* d_{A_0} u + *((d_{A_0} u) \wedge u^{-1} d_{A_0} u) = v,$$

where $v \in L^p(\mathfrak{g}_E)$ is the tautologically defined right-hand side and $u \in L^\infty \cap L_2^2$. Setting $b = d_{A_0} u$ and noting that $d_{A_0} b = F_{A_0} u$, with $F_{A_0} \in L_{k-1}^2(\Lambda^2 \otimes \mathfrak{g}_E) \subset L_1^2(\Lambda^2 \otimes \mathfrak{g}_E)$ and $F_{A_0} u \in L_1^2(\Lambda^2 \otimes \mathfrak{g}_E)$. Thus, we may conveniently rewrite (8.2) as a first-order elliptic equation in $b \in L_1^2(\Lambda^1 \otimes \mathfrak{g}_E)$,

$$(8.3) \quad (d_{A_0}^* + d_{A_0})b + *((*b) \wedge u^{-1}b) = v' \in L^p(\mathfrak{g}_E) \oplus L^p(\Lambda^2 \otimes \mathfrak{g}_E),$$

where $2 < p < 4$ and $v' = F_{A_0} u + v$. Finally, (8.3) can be rewritten as a local equation by writing $A_0 = \Gamma + a_0$, where Γ is the product connection in a local trivialization for E over a small ball $U \subset X$. Thus, the operator $d_{A_0}^* + d_{A_0}$ is replaced by $d^* + d$ in (8.2) and the additional terms are absorbed into the L^p inhomogeneous term v' to give:

$$(8.4) \quad (d^* + d)b + *((*b) \wedge u^{-1}b) = v'' \in L^p(U, \mathfrak{g}_E) \oplus L^p(U, \Lambda^2 \otimes \mathfrak{g}_E).$$

This is a first-order, elliptic equation with a quadratic non-linearity and Proposition 3.10 in [16] implies that the solution $b = d_{A_0} u \in L_1^2(U, \Lambda^1 \otimes \mathfrak{g}_E)$ is necessarily in $L_1^p(U', \Lambda^1 \otimes \mathfrak{g}_E)$ for $U' \Subset U$, provided $\|b\|_{L^4(U)} < \varepsilon(g, p, U)$, and so $u \in L_2^p(U', \mathfrak{g}_E)$. In particular, we find that $b \in L_1^p(X, \Lambda^1 \otimes \mathfrak{g}_E)$ and $u \in L_2^p(X, \mathfrak{g}_E)$ for any $2 < p < 4$, provided $\|d_{A_0} u\|_{L^4} < \varepsilon(g, p, X)$. The bootstrapping argument of Proposition 3.3 in [16] now implies that $d_{A_0} u \in L_k^2(X, \Lambda^1 \otimes \mathfrak{g}_E)$. Thus $u \in \mathcal{G}_E^{k+1}$ and $\chi \in L_{k+1}^2(X, \mathfrak{g}_E)$, as desired. \square

We can now proceed to the main argument:

Theorem 8.2. *Let X be a closed, Riemannian four-manifold and let G be compact Lie group. Then there are positive constants c, z with the following significance. Let E be a G bundle over X and suppose that $A_0 \in \mathcal{A}_E^2$, let $K_0[A_0] = (1 + \nu_0[A_0]^{-1})(1 + \|F_{A_0}\|_{L^2})$ and let ε_1 be a constant satisfying*

$$0 < \varepsilon_1 \leq z K_0^{-2}.$$

Then for any $A \in \mathcal{A}_E^2$ such that $\|A - A_0\|_{L_{1,A_0}^{1,2}} < \varepsilon_1$ there is a gauge transformation $u \in \mathcal{G}_E^3$ with the following properties:

- $d_{A_0}^*(u(A) - A_0) = 0$;
- $\|u(A) - A_0\|_{L_{1,A_0}^2} \leq cK_0\|A - A_0\|_{L_{1,A_0}^{\sharp,2}}$;
- $\|u - \text{id}_E\|_{L_{2,A_0}^{\sharp,2}} < cK_0\|A - A_0\|_{L_{1,A_0}^{\sharp,2}}$.

Proof. The argument is broadly similar to that of Lemma 3.6, except that we can show Ψ is a diffeomorphism directly — rather than just a local diffeomorphism — using the slightly stronger norms now at our disposal. Moreover, on this occasion we seek precise bounds on the solutions so we keep track of the dependence of constants on the curvature F_{A_0} and the least positive eigenvalue $\nu_0 = \nu_0[A_0]$ of the Laplacian $\Delta_{A_0} = d_{A_0}^* d_{A_0}$.

Write $A = A_0 + a$ and observe that

$$u(A) - A_0 = A - A_0 - (d_A u)u^{-1} = uau^{-1} - (d_{A_0} u)u^{-1}.$$

Recall that we have an L^2 -orthogonal decomposition

$$\Omega^0(\mathfrak{g}_E) = (\text{Ker } d_{A_0})^\perp \oplus \text{Ker } d_{A_0} = \text{Im } d_{A_0}^* \oplus \text{Ker } d_{A_0},$$

and that $d_{A_0}^* : L_1^2(\Lambda^1 \otimes \mathfrak{g}_E) \rightarrow L^2(\mathfrak{g}_E)$ has closed range; this gives

$$\begin{aligned} L_{2;A_0}^{\sharp,2}(\mathfrak{g}_E) &= (\text{Ker } d_{A_0}|_{L_{2;A_0}^{\sharp,2}})^\perp \oplus \text{Ker } d_{A_0}|_{L_{2;A_0}^{\sharp,2}} \\ &= (\text{Ker } d_{A_0}^*|_{L_{2;A_0}^{\sharp,2}})^\perp \oplus (\text{Im } d_{A_0}^*|_{L_{1,A_0}^{\sharp,2}}). \end{aligned}$$

We have a similar L^2 -orthogonal decomposition

$$\Omega^1(\mathfrak{g}_E) = \text{Im } d_{A_0} \oplus \text{Ker } d_{A_0}^* = (\text{Ker } d_{A_0}^*)^\perp \oplus \text{Ker } d_{A_0}^*,$$

and $d_{A_0} : L_1^2(\mathfrak{g}_E) \rightarrow L^2(\Lambda^1 \otimes \mathfrak{g}_E)$ has closed range; this leads to the L^2 -orthogonal decomposition

$$\begin{aligned} L_{1,A_0}^{\sharp,2}(\Lambda^1 \otimes \mathfrak{g}_E) &= (\text{Im } d_{A_0}|_{L_{2;A_0}^{\sharp,2}}) \oplus (\text{Ker } d_{A_0}^*|_{L_{1,A_0}^{\sharp,2}}) \\ &= (\text{Ker } d_{A_0}^*|_{L_{2;A_0}^{\sharp,2}})^\perp \oplus (\text{Ker } d_{A_0}^*|_{L_{1,A_0}^{\sharp,2}}). \end{aligned}$$

We now define a map

$$(8.5) \quad \begin{aligned} \Psi : (\text{Ker}(d_{A_0}|_{L_2^{\sharp,2}}))^\perp \oplus \text{Ker}(d_{A_0}^*|_{L_1^{\sharp,2}}) &\rightarrow L_{1,A_0}^{\sharp,2}(\Lambda^1 \otimes \mathfrak{g}_E), \\ (\chi, a) &\mapsto uau^{-1} - (d_{A_0} u)u^{-1}, \end{aligned}$$

where $u = e^\chi$ and the differential at (χ, a) given by

$$(8.6) \quad \begin{aligned} (D\Psi)_{(\chi,a)} : (\text{Ker}(d_{A_0}|_{L_2^{\sharp,2}}))^\perp \oplus \text{Ker}(d_{A_0}^*|_{L_1^{\sharp,2}}) &\rightarrow L_1^{\sharp,2}(\Lambda^1 \otimes \mathfrak{g}_E), \\ (\zeta, b) &\mapsto u(-d_A \oplus \iota)u^{-1}(\zeta, b) = u(-d_A \zeta + b)u^{-1}, \end{aligned}$$

since $(D\Psi)_{(0,a)}(\zeta, b) = -d_A \zeta + b$ and Ψ is \mathcal{G}_E -equivariant. Moreover, we have

$$(8.7) \quad (D^2\Psi)_{(\chi,a)}((\zeta, b), (\eta, \alpha)) = u[\eta, -d_A \zeta + b]u^{-1} + u[\alpha, \zeta]u^{-1}.$$

for $(\zeta, b), (\eta, \alpha) \in (\text{Ker}(d_{A_0}|_{L_2^{\sharp,2}}))^\perp \oplus \text{Ker}(d_{A_0}^*|_{L_1^{\sharp,2}})$.

We now verify that the conditions of the inverse function theorem (Theorem 3.2) hold for suitable constants K and δ . The operator

$$d_{A_0} : (\text{Ker}(d_{A_0}|_{L_2^{\sharp,2}}))^\perp \rightarrow (\text{Ker}(d_{A_0}^*|_{L_1^{\sharp,2}}))^\perp$$

has a two-sided inverse

$$G_{A_0}^0 d_{A_0}^* : (\text{Ker}(d_{A_0}^*|_{L_1^{\sharp,2}}))^{\perp} \rightarrow (\text{Ker}(d_{A_0}|_{L_2^{\sharp,2}}))^{\perp}.$$

Indeed, for $b \in (\text{Ker}(d_{A_0}^*|_{L_1^{\sharp,2}}))^{\perp}$, we have

$$\|G_{A_0}^0 d_{A_0}^* b\|_{L_{2,A_0}^{\sharp,2}} \leq c_0 K_0 \|d_{A_0}^* b\|_{L^{\sharp,2}} \leq c_0 K_0 \|b\|_{L_{1,A_0}^{\sharp,2}},$$

and so $G_{A_0}^0 d_{A_0}^*$ has $\text{Hom}(L_{1,A_0}^{\sharp,2}, L_{2,A_0}^{\sharp,2})$ operator norm bound

$$\|G_{A_0}^0 d_{A_0}^*\| \leq c_0 K_0.$$

In particular, we see that $(D\Psi)_{(0,0)}^{-1} = G_{A_0}^0 d_{A_0}^* \oplus \text{id}$ satisfies

$$(8.8) \quad \|(D\Psi)_{(0,0)}^{-1}\| \leq c_0 K_0$$

the first of the conditions we need to verify for $(D\Psi)_{(0,0)}$ in order to apply the inverse function theorem.

It remains to compare the differentials $(D\Psi)_{(\chi,a)}$ and $(D\Psi)_{(0,0)}$ using the mean value theorem,

$$(8.9) \quad (D\Psi)_{(\chi,a)}(\zeta, b) - (D\Psi)_{(0,0)}(\zeta, b) = \int_0^1 (D^2\Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a)) dt.$$

Thus, we need an estimate for $D^2\Psi$:

Claim 8.3. *There is a universal polynomial function $f(x, y)$, depending only on (X, g) and G , with $f(0, 0) = 0$, such that the following holds. For any $t \in [0, 1]$ we have:*

$$\begin{aligned} & \|(D^2\Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a))\|_{L_{1,A_0}^{\sharp,2}} \\ & \leq f(\|\chi\|_{L_{2,A_0}^{\sharp,2}}, \|a\|_{L_{1,A_0}^{\sharp,2}}) \left(\|\zeta\|_{L_{2,A_0}^{\sharp,2}} + \|b\|_{L_{1,A_0}^{\sharp,2}} \right). \end{aligned}$$

Proof. From (8.7) we have the $L^{\sharp,2}$ estimate

$$\begin{aligned} & \|(D^2\Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a))\|_{L^{\sharp,2}} \\ & \leq c \|\chi\|_{C^0} (\|d_{A_0}\zeta\|_{L^{\sharp,2}} + \|a\|_{L^{\sharp,2}} \|\zeta\|_{C^0} + \|b\|_{L^{\sharp,2}}) + c \|a\|_{L^{\sharp,2}} \|\zeta\|_{C^0}, \end{aligned}$$

and thus:

$$(8.10) \quad \begin{aligned} & \|(D^2\Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a))\|_{L^{\sharp,2}} \\ & \leq c \left(\|\chi\|_{L_{2,A_0}^{\sharp,2}} + \|a\|_{L^{\sharp,2}} \|\chi\|_{L_{2,A_0}^{\sharp,2}} + \|a\|_{L^{\sharp,2}} \right) \left(\|\zeta\|_{L_{2,A_0}^{\sharp,2}} + \|b\|_{L^{\sharp,2}} \right). \end{aligned}$$

The L^2 estimate of $\nabla_{A_0}(D^2\Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a))$ is given by

$$\begin{aligned} & \|\nabla_{A_0}(D^2\Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a))\|_{L^2} \\ & \leq c (\|\nabla_{A_0} u\|_{L^4} \|\chi\|_{C^0} + \|\nabla_{A_0} \chi\|_{L^4}) (\|d_{A_0}\zeta\|_{L^4} + \|a\|_{L^4} \|\zeta\|_{C^0} + \|b\|_{L^4}) \\ & \quad + c \|\chi\|_{C^0} (\|\nabla_{A_0}^2 \zeta\|_{L^2} + \|\nabla_{A_0} a\|_{L^2} \|\zeta\|_{C^0} + \|a\|_{L^4} \|\nabla_{A_0} \zeta\|_{L^4} + \|\nabla_{A_0} b\|_{L^2}) \\ & \quad + c \|\nabla_{A_0} u\|_{L^4} \|a\|_{L^4} \|\zeta\|_{L^4} + c \|\nabla_{A_0} a\|_{L^2} \|\zeta\|_{C^0} + c \|a\|_{L^4} \|\nabla_{A_0} \zeta\|_{L^4}, \end{aligned}$$

and hence, using Lemma 7.2 to estimate $u = e^\chi$ in terms of χ ,

$$(8.11) \quad \begin{aligned} & \|\nabla_{A_0}(D^2\Psi)_{(t_\chi, ta)}((\zeta, b), (\chi, a))\|_{L^2} \\ & \leq f_1\left(\|\chi\|_{L_{2,A_0}^{\sharp,2}}, \|a\|_{L_{1,A_0}^2}\right) \left(\|\zeta\|_{L_{2,A_0}^{\sharp,2}} + \|b\|_{L_{1,A_0}^2}\right), \end{aligned}$$

where $f_1(x, y)$ is a polynomial function with $f_1(0, 0) = 0$.

Noting that $d_{A_0}^* a = 0$, we have

$$(8.12) \quad \begin{aligned} d_{A_0}^*[a, \zeta] &= d_{A_0}^*(a\zeta - \zeta a) \\ &= (d_{A_0}^* a)\zeta - a \wedge d_{A_0} \zeta - *(d_{A_0} \zeta \wedge *a) - \zeta(d_{A_0}^* a) \\ &= -a \wedge d_{A_0} \zeta - *(d_{A_0} \zeta \wedge *a). \end{aligned}$$

and similarly for $[\chi, b]$ since $d_{A_0}^* b = 0$. For any $\beta \in L_1^2(\Lambda^1 \otimes \mathfrak{g}_E)$ we have

$$(8.13) \quad \begin{aligned} d_{A_0}^*(u\beta u^{-1}) &= - * d_{A_0}(u(*\beta)u^{-1}) \\ &= - * (d_{A_0} u \wedge *\beta u^{-1}) + u(d_{A_0}^* \beta)u^{-1} - *u((*\beta) \wedge u(d_{A_0} u)u^{-1}). \end{aligned}$$

Therefore, equations (8.7), (8.12), and (8.13) and the estimates for $u = e^\chi$ in Lemma 7.2 yield

$$(8.14) \quad \begin{aligned} & \|d_{A_0}^*(D^2\Psi)_{(\chi, a)}((\zeta, b), (\chi, a))\|_{L^{\sharp,2}} \\ & \leq \|d_{A_0}^*(u[\chi, -d_A \zeta + b]u^{-1} + u[a, \zeta]u^{-1})\|_{L^{\sharp,2}} \\ & \leq f_2(\|\chi\|_{L_{2,A_0}^{\sharp,2}}, \|a\|_{L_{2,A_0}^{\sharp,4}}) \left(\|\zeta\|_{L_{2,A_0}^{\sharp,2}} + \|b\|_{L_{2,A_0}^{\sharp,4}}\right), \end{aligned}$$

where $f_2(x, y)$ is a polynomial function with $f_2(0, 0) = 0$. The claim now follows by combining (8.10), (8.11), and (8.14). \square

Therefore, from Claim 8.3 and (8.9) we have

$$(8.15) \quad \begin{aligned} & \|(D\Psi)_{(\chi, a)}(\zeta, b) - (D\Psi)_{(0,0)}(\zeta, b)\|_{L_{1,A_0}^{\sharp,2}} \\ & \leq f(\|\chi\|_{L_{2,A_0}^{\sharp,2}}, \|a\|_{L_{1,A_0}^{\sharp,2}}) \left(\|\zeta\|_{L_{2,A_0}^{\sharp,2}} + \|b\|_{L_{1,A_0}^{\sharp,2}}\right). \end{aligned}$$

Consequently, with respect to the $\text{Hom}(L_{2,A_0}^{\sharp,2}, L_{1,A_0}^{\sharp,2})$ operator norm, (8.15) yields the bound

$$(8.16) \quad \|(D\Psi)_{(\chi, a)} - (D\Psi)_{(0,0)}\| \leq \frac{1}{2}c_0^{-1}K_0^{-1},$$

where $c_0 K_0 = K$ is the constant of (8.8), provided (χ, a) satisfies the constraint

$$(8.17) \quad \|\chi\|_{L_{2,A_0}^{\sharp,2}} + \|a\|_{L_{1,A_0}^{\sharp,2}} \leq c_1 K_0^{-1} = \delta.$$

Define balls centered at the origins in $(\text{Ker}(d_{A_0}|_{L_{2,A_0}^{\sharp,2}}))^{\perp}$ and $\text{Ker}(d_{A_0}^*|_{L_{1,A_0}^{\sharp,2}})$ by setting

$$\begin{aligned} B_0^{\perp;2,\sharp,2}(\delta) &= \{\chi \in (\text{Ker}(d_{A_0}|_{L_{2,A_0}^{\sharp,2}}))^{\perp} : \|\chi\|_{L_{2,A_0}^{\sharp,2}} < \delta\}, \\ \mathbf{B}_0^{1,\sharp,2}(\delta) &= \{a \in \text{Ker}(d_{A_0}^*|_{L_{1,A_0}^{\sharp,2}}) : \|a\|_{L_{1,A_0}^{\sharp,2}} < \delta\}. \end{aligned}$$

Hence, Theorem 3.2 implies that the map

$$\Psi : B_0^{\perp;2,\sharp,2}(\delta) \times \mathbf{B}_0^{1,\sharp,2}(\delta) \rightarrow \mathcal{A}_E^{1,\sharp,2}$$

is injective, its image is an open subset of $\mathcal{A}_E^{1,\sharp,2}$ and contains the ball $B_{A_0}^{1,\sharp,2}(\delta/(2K))$, the inverse map Ψ^{-1} is a diffeomorphism from $B_{A_0}^{1,\sharp,2}(\delta/(2K))$ onto its image, and if (χ_1, A_1) , (χ_2, A_2) are points in $B_0^{\perp,2,\sharp,2}(\delta) \times \mathbf{B}_0^{1,\sharp,2}(\delta)$, then

$$\|\chi_1 - \chi_2\|_{L_{2,A_0}^{\sharp,2}} + \|A_1 - A_2\|_{L_{1,A_0}^{\sharp,2}} \leq 2K\|u_1(A_1) - u_2(A_2)\|_{L_{1,A_0}^{\sharp,2}},$$

where $u_i = e^{\chi_i}$, $i = 1, 2$. In particular, setting $(\chi_2, A_2 - A_0) = (0, 0)$, we see that if A is a point in $\mathcal{A}_E^{1,\sharp,2}$ such that $\|A - A_0\|_{L_{1,A_0}^{\sharp,2}} < \delta/(2K)$, then there is a unique solution $(\chi, u^{-1}(A)) = \Psi^{-1}(A)$ in $B_0^{\perp,2,\sharp,2}(\delta) \times \mathbf{B}_0^{1,\sharp,2}(\delta)$. Here, $u = e^\chi$ is a gauge transformation with $\chi \in B_0^{\perp,2,\sharp,2}(\delta)$ such that

$$(8.18) \quad d_{A_0}^*(u^{-1}(A) - A_0) = 0,$$

$$\|\chi\|_{L_{2,A_0}^{\sharp,2}} + \|u^{-1}(A) - A_0\|_{L_{1,A_0}^{\sharp,2}} \leq 2K\|A - A_0\|_{L_{1,A_0}^{\sharp,2}}.$$

Lemma 7.2 implies that $u = e^\chi$ satisfies

$$(8.19) \quad \|u - \text{id}_E\|_{L_{2,A_0}^{\sharp,2}} \leq f_3(\|\chi\|_{L_{2,A_0}^{\sharp,2}}) \leq c\|\chi\|_{L_{2,A_0}^{\sharp,2}} \leq c_2\delta,$$

where $f_3(x)$ is a polynomial with coefficients depending only on (X, g) and G such that $f_3(0) = 0$. Noting that $K = c_0K_0$, $\delta = c_1K_0^{-1}$, and $\delta/(2K) = \frac{1}{2}c_0c_1K_0^{-2}$, the desired estimates follows from (8.18) and (8.19). Finally, Lemma 8.1 implies that $u \in \mathcal{G}_E^3$ and this completes the proof of the theorem. \square

While the L_1^2 estimate of Theorem 8.2 suffices for most practical purposes, it is occasionally useful to have the slightly weaker $L^{2,\sharp,4}$ bound at hand. Recall from Section 4 that we defined

$$\|a\|_{\mathcal{L}_{1,A_0}^{\sharp,2}} = \|a\|_{L^{2,\sharp,4}} + \|d_{A_0}^*a\|_{L^{\sharp,2}}, \quad a \in \Omega^1(\mathfrak{g}_E).$$

A slight modification of the proof of Theorem 8.2 yields:

Theorem 8.4. *Continue the hypotheses of Theorem 8.2. Then for any $A \in \mathcal{A}_E^2$ such that $\|A - A_0\|_{\mathcal{L}_{1,A_0}^{\sharp,2}} < \varepsilon_1$ there is a gauge transformation $u \in \mathcal{G}_E^3$ with the following properties:*

- $d_{A_0}^*(u(A) - A_0) = 0$;
- $\|u(A) - A_0\|_{L^{2,\sharp,4}} \leq cK_0\|A - A_0\|_{\mathcal{L}_{1,A_0}^{\sharp,2}}$;
- $\|u - \text{id}_E\|_{L_{2,A_0}^{\sharp,2}} < cK_0\|A - A_0\|_{\mathcal{L}_{1,A_0}^{\sharp,2}}$.

Proof. The first difference in the argument is that the map Ψ in (8.5) is replaced by

$$(8.20) \quad \Psi : (\text{Ker}(d_{A_0}|_{L_2^{\sharp,2}}))^{\perp} \oplus \text{Ker}(d_{A_0}^*|_{\mathcal{L}_{1,A_0}^{\sharp,2}}) \rightarrow \mathcal{L}_{1,A_0}^{\sharp,2}(\Lambda^1 \otimes \mathfrak{g}_E),$$

$$(\chi, a) \mapsto uau^{-1} - (d_{A_0}u)u^{-1}.$$

The second difference is that Claim 8.3 is replaced by:

Claim 8.5. *There is a universal polynomial function $f(x, y)$, depending only on (X, g) and G , with $f(0, 0) = 0$, such that the following holds. For any $t \in [0, 1]$ we have:*

$$\|(D^2\Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a))\|_{\mathcal{L}_{1,A_0}^{\sharp,2}} \leq f(\|\chi\|_{L_{2,A_0}^{\sharp,2}}, \|a\|_{L^{2,\sharp,4}}) \left(\|\zeta\|_{L_{2,A_0}^{\sharp,2}} + \|b\|_{L^{2,\sharp,4}} \right).$$

Proof. From (8.7) we now have the $L^{2\sharp,4}$ estimate

$$\begin{aligned} & \| (D^2 \Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a)) \|_{L^{2\sharp,4}} \\ & \leq c \|\chi\|_{C^0} (\|d_{A_0} \zeta\|_{L^{2\sharp,4}} + \|a\|_{L^{2\sharp,4}} \|\zeta\|_{C^0} + \|b\|_{L^{2\sharp,4}}) + c \|a\|_{L^{2\sharp,4}} \|\zeta\|_{C^0}, \end{aligned}$$

and thus:

$$\begin{aligned} (8.21) \quad & \| (D^2 \Psi)_{(t\chi, ta)}((\zeta, b), (\chi, a)) \|_{L^{2\sharp,4}} \\ & \leq c \left(\|\chi\|_{L^{2\sharp,2}_{2,A_0}} + \|a\|_{L^{2\sharp,4}} \|\chi\|_{L^{2\sharp,2}_{2,A_0}} + \|a\|_{L^{2\sharp,4}} \right) \left(\|\zeta\|_{L^{2\sharp,2}_{2,A_0}} + \|b\|_{L^{2\sharp,4}} \right). \end{aligned}$$

Combining (8.14) and (8.21) yields the claim. \square

The rest of the argument proceeds exactly as before and completes the proof of the theorem. \square

We now have our second proof of Theorem 6.1 via Theorems 8.2 and 8.4:

Proof of Theorem 6.1. From the hypotheses we have $A_0 \in \mathcal{A}_E^k$ and $[A] \in \mathcal{B}_E^k$ with $k \geq 2$. According to Lemma 6.3, there is gauge transformation $w \in \mathcal{G}_E^3$ such that

$$\text{dist}_{L^{2\sharp,2}_{1,A_0}}([A], [A_0]) = \|w(A) - A_0\|_{L^{2\sharp,2}_{1,A_0}},$$

where $A \in \mathcal{A}_E^k$, so Theorems 8.2 and the argument of 8.4 imply that there is a gauge transformation $v \in \mathcal{G}_E^3$ so that $u(A)$ satisfies the conclusions of Assertion (2) with $u = vw \in \mathcal{G}_E^3$. Since $d_{A_0}^*(u(A) - A_0) = 0$ and $u \in \mathcal{G}_E^3$ and $A, A_0 \in \mathcal{A}_E^k$, a standard bootstrapping argument implies that $u \in \mathcal{G}_E^{k+1}$.

Similarly, by Lemma 6.3, there is gauge transformation $w \in \mathcal{G}_E^3$ such that

$$\text{dist}_{L^{2\sharp,2}_{1,A_0}}([A], [A_0]) = \|w(A) - A_0\|_{L^{2\sharp,2}_{1,A_0}},$$

so Assertion (1) follows from Theorem 8.4 in the same manner. \square

REFERENCES

- [1] R. A. Adams, *Sobolev Spaces*, Academic Press, Orlando, FL, 1975.
- [2] M. F. Atiyah, N. J. Hitchin, and I. M. Singer, *Self-duality in four-dimensional Riemannian geometry*, Proc. Royal Soc. London **A 362** (1978) 425–461.
- [3] T. Aubin, *Nonlinear Analysis on Complex Manifolds. Monge-Ampère Equations*, Springer, New York, 1982.
- [4] S. B. Bradlow, *Special metrics and stability for holomorphic bundles with global sections*, J. Differential Geom. **33** (1991) 169–213.
- [5] S. B. Bradlow and G. D. Daskalopoulos, *Moduli of stable pairs for holomorphic bundles over Riemann surfaces*, Internat. J. Math. **2** (1991) 477–513.
- [6] G. Daskalopoulos, private communication.
- [7] S. K. Donaldson, *A new proof of a theorem of Narasimhan and Seshadri*, J. Differential Geom. **18** (1983) 269–277.
- [8] ———, *Anti-self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles*, Proc. London Math. Soc. **50** (1985) 1–26.
- [9] ———, *Connections, cohomology and the intersection forms of four-manifolds*, J. Differential Geom. **24** (1986) 275–341.
- [10] ———, *Irrationality and the h-cobordism conjecture*, J. Differential Geom. **26** (1987), 141–168.
- [11] ———, *Polynomial invariants for smooth 4-manifolds*, Topology **29**, 257–315 (1990).
- [12] ———, *The approximation of instantons*, Geom. Funct. Anal. **3** (1993) 179–200.
- [13] ———, *The Seiberg-Witten equations and four-manifold topology*, Bull. Amer. Math. Soc. **33** (1996) 45–70.

- [14] S. K. Donaldson and P. B. Kronheimer, *The Geometry of Four-Manifolds*, Oxford University Press, Oxford, 1990.
- [15] S. K. Donaldson and D. P. Sullivan, *Quasi-conformal four-manifolds*, Acta Math. **163** (1990) 181–252.
- [16] P. M. N. Feehan and T. G. Leness, *PU(2) monopoles. I: Regularity, Uhlenbeck compactness, and transversality*, J. Differential Geom. **49** (1998), 265–410, [dg-ga/9710032](#).
- [17] ———, *PU(2) monopoles and relations between four-manifold invariants*, Topology Appl. **88** (1998), 111–145, [dg-ga/9709022](#).
- [18] ———, *PU(2) monopoles, II: Highest-level singularities and relations between four-manifold invariants*, submitted to a print journal, December 8, 1997; [dg-ga/9712005](#).
- [19] ———, *PU(2) monopoles, III: Existence of gluing and obstruction maps*, in preparation.
- [20] ———, *PU(2) monopoles, IV: Surjectivity of gluing maps*, in preparation.
- [21] ———, *in preparation*.
- [22] D. Freed and K. K. Uhlenbeck, *Instantons and Four-Manifolds*, 2nd ed., Springer, New York, 1991.
- [23] R. Friedman and J. W. Morgan, *Smooth Four-manifolds and Complex Surfaces*, Springer, Berlin, 1994.
- [24] D. Gilbarg and N. Trudinger, *Elliptic Partial Differential Equations of Second Order*, 2nd ed., Springer, New York, 1983.
- [25] L. Göttsche, *Modular forms and Donaldson invariants for 4-manifolds with $b_+ = 1$* , J. Amer. Math. Soc. **9** (1996), 827–843; [alg-geom/9506018](#).
- [26] D. Kotschick and J. W. Morgan, *SO(3) invariants for four-manifolds with $b^+ = 1$. II*, J. Differential Geom. **39** (1994) 433–456.
- [27] P. B. Kronheimer and T. S. Mrowka, *Embedded surfaces and the structure of Donaldson’s polynomial invariants*, J. Differential Geom. **43** (1995) 573–734.
- [28] H. B. Lawson, *The Theory of Gauge Fields in Four Dimensions*, Conf. Board Math. Sci. **58**, Amer. Math. Soc., Providence, RI, 1985.
- [29] J. Li, *Algebraic geometric interpretation of Donaldson’s polynomial invariants*, J. Differential Geom. **37** (1993) 417–466.
- [30] H. B. Lawson and M-L. Michelsohn, *Spin Geometry*, Princeton Univ. Press, Princeton, NJ, 1988.
- [31] P. K. Mitter and C. M. Viallet, *On the bundle of connections and the gauge orbit manifold in Yang-Mills theory*, Comm. Math. Phys. **79** (1981) 457–472.
- [32] J. W. Morgan, *Gauge Theory and the Topology of Smooth Four-Manifolds*, Lecture Notes, Harvard University, 1988.
- [33] ———, *Comparison of the Donaldson polynomial invariants with their algebro-geometric analogues*, Topology **32** (1993) 449–488.
- [34] J. W. Morgan and P. S. Ozsváth, private communication.
- [35] C. Okonek and A. Teleman, *Quaternionic monopoles*, Comm. Math. Phys. **180** (1996) 363–388; [alg-geom/9505029](#).
- [36] R. S. Palais, *Ljusternik-Snirelman theory on Banach manifolds*, Topology **5** (1966) 115–132.
- [37] ———, *Foundations of Global Non-Linear Analysis*, Benjamin, New York, 1968.
- [38] ———, *Critical-point theory and the min-max principle*, Proc. Symp. Pure Math. **15**, Amer. Math. Soc., Providence, RI, 1970.
- [39] T. H. Parker, *Gauge theories on four dimensional Riemannian manifolds*, Comm. Math. Phys. **85** (1982) 563–602.
- [40] T. H. Parker and C. H. Taubes, *On Witten’s proof of the positive energy theorem*, Comm. Math. Phys. **84** (1982) 223–238.
- [41] V. Y. Pidstrigach, Lecture at the Newton Institute, Cambridge, December 1994.
- [42] V. Y. Pidstrigach and A. N. Tyurin, *Localization of Donaldson invariants along the Seiberg-Witten classes*, [dg-ga/9507004](#).
- [43] I. M. Singer, *Some remarks on the Gribov ambiguity*, Comm. Math. Phys. **60** (1978) 7–12.
- [44] E. Stein, *Singular integral operators and differentiability properties of functions*, Princeton University Press, Princeton, NJ, 1970.
- [45] C. H. Taubes, *Self-dual Yang-Mills connections on non-self-dual 4-manifolds*, J. Differential Geom. **17** (1982) 139–170.
- [46] ———, *Stability in Yang-Mills theories*, Comm. Math. Phys. **91** (1983) 235–263.
- [47] ———, *Self-dual connections on 4-manifolds with indefinite intersection matrix*, J. Differential Geom. **19** (1984) 517–560.

- [48] ———, *Path-connected Yang-Mills moduli spaces*, J. Differential Geom. **19** (1984) 337–392.
- [49] ———, *A framework for Morse theory for the Yang-Mills functional*, Invent. Math. **94** (1988) 327–402.
- [50] ———, *The stable topology of self-dual moduli spaces*, J. Differential Geom. **29** (1989) 162–230.
- [51] ———, *The existence of anti-self-dual conformal structures*, J. Differential Geom. **36** (1992) 163–253.
- [52] ———, *Metrics, Connections, and Gluing Theorems*, Conf. Board Math. Sci. **89**, Amer. Math. Soc., Providence, RI, 1996.
- [53] A. Teleman, *Non-abelian Seiberg-Witten theory and stable oriented pairs*, Internat. J. Math. **8** (1997) 507–535; `alg-geom/9609020`.
- [54] K. K. Uhlenbeck, *Connections with L^p bounds on curvature*, Comm. Math. Phys. **83** (1982) 31–42.
- [55] E. Witten, *Monopoles and four-manifolds*, Math. Res. Lett. **1** (1994) 769–796.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, ONE OXFORD STREET, CAMBRIDGE, MA 02138

E-mail address: `feehan@math.harvard.edu`